arXiv:0904.4834v2 [math.AG] 1 Dec 2009

GROMOV-WITTEN GAUGE THEORY I

EDWARD FRENKEL, CONSTANTIN TELEMAN, AND A.J. TOLLAND

Abstract. We introduce a modular completion of the stack of maps from stable marked curves to the quotient stack $[\mathrm{pt}/\mathbb{C}^\times]$, and use this stack to construct some gauge-theoretic analogues of the Gromov-Witten invariants. We also indicate the generalization of these invariants to the quotient stacks $[X/\mathbb{C}^\times]$, where X is a smooth proper complex algebraic variety.

Contents

1.	The Stack of Gieseker Bundles	6
2.	The Geometry of $\widetilde{\mathfrak{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$	15
3.	The Structure of the Local Atlas	19
4.	Admissible Classes	27
5.	Gromov-Witten Invariants for $[\mathrm{pt}/\mathbb{C}^\times]$	30
6.	Towards Gromov-Witten Invariants for $[X/\mathbb{C}^\times]$	33
References		39

In this paper we construct algebro-geometric Gromov-Witten invariants for the quotient stack $[\mathrm{pt}/\mathbb{C}^\times]$. These invariants are the "twisted" indices of products of evaluation K-theory classes on certain moduli stacks $\widetilde{\mathfrak{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$, which classify maps from marked nodal curves to the quotient stack $[\mathrm{pt}/\mathbb{C}^\times]$. These are the moduli stacks of principal $\mathbb{C}^\times$-bundles on such curves. We do not impose stability conditions on these bundles, and our moduli stacks are Artin stacks that are very far from proper. Consequently, the existence of these invariants is non-trivial. Our main theorem asserts that they are, in fact, well-defined.

This construction is the first step in a larger project, the goal of which is to define Gromov-Witten invariants for the Artin stacks $[X/G]$, where X is a smooth projective variety and G is a reductive algebraic group over $\mathbb{C}$. Similar invariants (in the case where G is a finite group) have already been defined in the symplectic setting by Chen & Ruan [CR02] and algebro-geometrically by Abramovich, Graber, Olsson, & Vistoli [AV02, AGV08]. In this paper, we explain (modulo a certain conjecture) how to define invariants for $[X/\mathbb{C}^\times]$ in terms of our invariants for $[\mathrm{pt}/\mathbb{C}^\times]$.

Partially supported by DARPA and AFOSR through the grant FA9550-07-1-0543.

We expect that these invariants – defined as twisted indices of products of evaluation classes on moduli stacks of maps from marked curves to $[X/\mathbb{C}^\times]$ – may be interpreted as correlation functions in a topological quantum field theory. This quantum field theory is in some sense a *gauge* theory. The stack of algebraic principal G-bundles on a smooth curve Σ is homotopy equivalent [AB83] to the stack of K-connections on Σ, where K is the compact form of G, and so one can view our construction of Gromov-Witten invariants for $[\mathrm{pt}/\mathbb{C}^\times]$ as a topological version of Feynman path integration over the space of $U(1)$-connections.

Similar invariants have been studied by Mundet, Cieliebak, Gaio, Salamon, and Tian [MiR00, CGS00, CGMiRS02, MiR03, MiRT04], who defined invariants for symplectic manifolds with Hamiltonian $U(1)$-actions by integrating over coarse moduli spaces of solutions to the symplectic vortex equations, and by Gonzalez & Woodward [GW08, GW09], who considered the generalization from $U(1)$ to compact reductive groups. In fact, the idea that the path integral of a gauged nonlinear sigma model might compute cohomological invariants of moduli spaces of connections A and $\overline{\partial}_A$-holomorphic sections of a bundle with fiber X appears in Witten's original paper on topological sigma models [Wit88]. For a more recent perspective on this, see the study [FLN08] of gauged sigma models in the "infinite radius limit".

Our invariants differ from the existing ones in that we do not impose stability conditions in the sense of geometric invariant theory on the maps to $[X/\mathbb{C}^\times]$ and the invariants take values in K-theory rather than cohomology. (Our approach has been outlined, in particular, in a lecture given by one of us at the Seattle conference in 2005 [Tel05].) The proof of finiteness of these invariants uses a technical argument adapted from [Tel00]. Consequently, our invariants are defined in greater generality than the known ones. Indeed, we expect to recover the Gromov-Witten invariants of GIT quotients from our invariants by applying the Chern character to certain limits of our invariants. This was done for smooth curves and maps to $[\mathrm{pt}/G]$ in [TW03].

0.1. Sketch of the Construction.

We explain here the construction of invariants for $[\mathrm{pt}/\mathbb{C}^\times]$. (The invariants for $[X/\mathbb{C}^\times]$ are built up from this special case.)

To begin, we need a completion of the stack $\mathrm{Bun}_{\mathbb{C}^\times}(g, I)$ of maps from stable marked curves to $[\mathrm{pt}/\mathbb{C}^\times]$. Recall that defining a $\mathbb{C}^\times$-bundle on a nodal curve Σ is the same as defining a $\mathbb{C}^\times$-bundle on the normalization of Σ together with the data of identifications of the fibers at the preimages of the nodal points. $\mathrm{Bun}_{\mathbb{C}^\times}(g, I)$ fails to be complete because the space of identifications over a given node is isomorphic to $\mathbb{C}^\times$. Following Gieseker [Gie84] and Caporaso [Cap94], we get a completion – denoted $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ – by adding new strata which represent the limits where an identification goes to zero or infinity by allowing projective lines carrying the line bundle $\mathcal{O}_{\mathbb{P}^1}(1)$ to appear at the nodes. (Similar completions of various stacks of vector bundles on nodal curves have been studied by many authors; see the reviews [Kle84, Ale04]. Our definition was inspired by Caporaso's thesis [Cap94] and the papers of Nagaraj & Seshadri [NS97, NS99].)

The stack $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ has a forgetful map

$$F : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to \overline{\mathcal{M}}_{g,I}.$$

It also carries a) a universal curve $\pi : C \to \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ with marked points $\sigma_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to C$ and b) a universal principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$, representing the universal map $\phi : C \to [\mathrm{pt}/\mathbb{C}^\times]$. Composition gives the evaluation maps $\mathrm{ev}_i = \phi \circ \sigma_i$.

$$\mathrm{ev}_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to [\mathrm{pt}/\mathbb{C}^\times].$$

A vector bundle on $[\mathrm{pt}/\mathbb{C}^\times]$ is a $\mathbb{C}^\times$-representation V. Pulling these back along the universal morphism ϕ gives the universal vector bundles $\mathcal{V} = \phi^* V$ on C. Likewise, if we pull K-theory classes $[V] \in K([\mathrm{pt}/\mathbb{C}^\times]) \simeq K_{\mathbb{C}^\times}(\mathrm{pt})$ back along the maps ev_i, we obtain the *evaluation classes* $\mathrm{ev}_i^*[V]$.

The Gromov-Witten invariants of $[\mathrm{pt}/\mathbb{C}^\times]$ (like those of a variety X) result from pushing a product of evaluation classes forward along the forgetful morphism F. However, our setup differs from the standard one in two ways.

1. Our invariants are constructed in K-theory, rather than cohomology.
2. Our invariants are always *twisted*, in the sense of [CG07].

Twisting requires some definition. A line bundle $\mathcal{L}$ on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is said to be *admissible* if (topologically)

$$\mathcal{L}^{-1} \simeq (\det R\pi_* \phi^* V)^{\otimes r},$$

where V is a non-trivial irreducible $\mathbb{C}^\times$-representation and r is a positive rational number. An *$\mathcal{L}$-twisted Gromov-Witten invariant of* $[\mathrm{pt}/\mathbb{C}^\times]$ is the Euler characteristic (on $\overline{\mathcal{M}}_{g,I}$) of the twist by (admissible) $\mathcal{L}$ of a product of evaluation classes:

$$\langle V_1, ..., V_{|I|} \rangle_{\mathcal{L}} = \chi_{\overline{\mathcal{M}}_{g,I}} \Big(RF_* \big(\mathcal{L} \bigotimes \otimes_{i \in I} \mathrm{ev}_i^* V_i \big) \Big).$$

One can also twist by powers of the index classes $R\pi_* \mathcal{V}$ of vector bundles $\mathcal{V}$ on C; these classes may be assembled into *higher twistings*. An *admissible complex* is a sum of products of complexes of the form

$$\mathcal{L} \bigotimes \otimes_a (R\pi_* \mathcal{V}_a)^{n_a} \bigotimes \otimes_i (\mathrm{ev}_i^* V_i \otimes T_i^{\otimes n_i}).$$

The subring of $K(\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ generated by such products is called the *ring of admissible classes*. It is a subring without unit; the trivial line bundle $\mathcal{O}$ is not admissible. It is not obvious that the push-forward of an admissible class along F is well-defined, because the moduli stacks $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ differ from Kontsevich's stack of stable maps in two important ways.

First, $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is an Artin stack, rather than a Deligne-Mumford stack; points can have a finite-dimensional stabilizers. It is for this reason that we use

K-theory instead of cohomology. The forgetful morphism F to the stack of curves is always Artin, and it is generally impossible to push cohomology classes forward along such morphisms. The problem can be understood as follows. The fibers of F are quotients stacks of the form $[A/\mathcal{G}]$ ($\mathcal{G}$ a group), and integrating over $[A/\mathcal{G}]$ amounts to first pushing forward along $[A/\mathcal{G}] \to [\mathrm{pt}/\mathcal{G}]$ and then pushing forward along $[\mathrm{pt}/\mathcal{G}] \to \mathrm{pt}$. But the latter pushforward is zero in cohomology. Indeed, pushing forward along $[\mathrm{pt}/\mathcal{G}] \to \mathrm{pt}$ shifts the degree of a cohomology class by $-\dim([\mathrm{pt}/\mathcal{G}]) = \dim(\mathcal{G})$, so the only classes with non-zero push-forward live in $H_{\mathcal{G}}^{-\dim(\mathcal{G})}(\mathrm{pt})$. But $H^n_{\mathcal{G}}(\mathrm{pt})$ vanishes if n is negative. In K-theory, on the other hand, the pushforward along $[\mathrm{pt}/\mathcal{G}] \to \mathrm{pt}$ does exist. The class $[V] \in K_{\mathcal{G}}(\mathrm{pt})$ represented by a $\mathcal{G}$-module V is sent to the K-theory class $[V^{\mathcal{G}}] \in K(\mathrm{pt})$ represented by the vector space of $\mathcal{G}$-invariants in V.

The second problem is that $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is very far from proper. It is complete, but it is in general neither separated nor of finite type. Thus, the existence of a pushforward along the forgetful morphism to $\overline{\mathcal{M}}_{g,I}$ is a delicate matter; not every K-theory class on $\overline{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ has a well-defined index. The main theorem in this paper asserts that the particular classes we want to push forward do indeed have well-defined index.

(There is a third difference, which makes things easier: The stack $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is unobstructed, so we don't need to use virtual structure sheaves.)

Main Theorem. *The derived pushforward $RF_*\alpha$ of an admissible complex α along the bundle-forgetting map $F : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to \overline{\mathcal{M}}_{g,I}$ is a complex of coherent sheaves.*

This theorem is a relative version (allowing the curve to vary) of the finiteness theorem proved on $\mathrm{Bun}_G(\Sigma)$ in [TW03]. The proof, in rough outline:

1. Coherence is a local property, so we can work in an affine étale neighborhood B in $\overline{\mathcal{M}}_{g,I}$. For small enough B, the restriction of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ to B can be presented as a stack quotient $[A/\mathcal{G}]$, where A is an algebraic space classifying $\mathbb{C}^\times$-bundles equipped with trivializations at marked points labelled by a set V, and $\mathcal{G} = (\mathbb{C}^\times)^V$. Since B is affine, we can prove coherence by showing that the $\mathcal{G}$-invariants in the global sections $R\Gamma(A, \alpha)$ are finitely-generated over B.
2. The Kirwan-Ness stratification associated to the $\mathcal{G}$-action on A is compatible with the stratification by degeneracy type, which tracks the multidegrees of the bundles. This fact makes it possible to identify algebraic subspaces $S \subset A$ for which $[S/\mathcal{G}] \simeq Q \times [\mathrm{pt}/\mathbb{C}^\times]$, with Q proper over B.
3. The $\mathcal{G}$-weights of the fixed point fibers of admissible line bundles grow linearly with the multidegree, while the weights of evaluation classes and their descendants and index classes are bounded functions of the multidegree. These estimates make it possible to prove that the $\mathcal{G}$-invariants in

the local cohomologies of admissible complexes on the degeneracy strata are always coherent, and are non-zero in only finitely many cases. This result allows us to reduce the question of finite-generation of invariants on A to S, where it follows trivially from the properness result.

0.2. **Invariants for** $[X/\mathbb{C}^\times]$**.** Recall that $[X/\mathbb{C}^\times]$ is defined so that maps from a curve Σ to $[X/\mathbb{C}^\times]$ correspond to pairs $(\mathcal{P}, s)$ consisting of a principal $\mathbb{C}^\times$-bundle and a section $s \in \Gamma(\Sigma, \mathcal{P} \times_{\mathbb{C}^\times} X)$ of the associated bundle with fiber X. To define Gromov-Witten invariants for $[X/\mathbb{C}^\times]$, we need a moduli stack $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$ of curves and degree β maps to $[X/\mathbb{C}^\times]$ on which we can define tautological classes, and we need a way of defining the pushforward of these classes from $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$ to $\overline{\mathcal{M}}_{g,I}$.

In this paper, we define an appropriate moduli stack $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$. This stack has a natural section-forgetting morphism

$$F_\beta : \widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times]) \to \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]).$$

The fibers of this morphism are stacks of sections of bundles with fiber X associated to Gieseker bundles. Such sections are locally maps to X, so they can develop singularities in the same way. Following Kontsevich, we ensure that F_β is proper by allowing bubbling at points where such singularities occur.

Thus, the morphism F_β is very much like the morphism $F_\beta : \overline{\mathcal{M}}_{g,I,\beta}(X) \to \overline{\mathcal{M}}_{g,I}$ used in ordinary Gromov-Witten theory. We prove, in fact, that the morphism F_s is proper, Deligne-Mumford, and carries a perfect obstruction theory, relative to $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$, which is smooth. (The proofs are straightforward generalizations of the usual ones in Gromov-Witten theory.) These facts imply the existence of a virtual K-theoretic pushforward along F_β. We conjecture that the virtual pushforward of an admissible class along F_β is an admissible class on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$. If this conjecture holds, then we can safely define the Gromov-Witten invariants of $[X/\mathbb{C}^\times]$ to be the K-theory classes on $\overline{\mathcal{M}}_{g,I}$ obtained by applying $F_{\beta*}^{vir}$ and $F_{\mathcal{P}*}$ to an admissible class on $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$.

0.3. **Plan of the Paper.** In Section 1, we review basic facts about principal $\mathbb{C}^\times$-bundles on curves, then define the moduli stack $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ of Gieseker bundles on stable curves, and discuss a few examples (small g and $|I|$).

In Section 2, we introduce an atlas A for our stack, and discuss the deformations of Gieseker bundles. We then draw a few basic conclusions about the geometry of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$.

In Section 3, we show that the Kirwan-Ness stratification of A (for its given $\mathcal{G} = (\mathbb{C}^\times)^V$-action) is compatible with (a certain refinement of) the induced degeneracy stratification. We also introduce some finite-type subspaces of A and explain how to think of them in terms of the stratification.

In Section 4, we define admissible classes and estimate the behavior of the $\mathcal{G}$-weights of their fixed point fibers as functions of the multidegrees of the bundles classified by the fixed points.

In Section 5, we define the Gromov-Witten invariants for $[\mathrm{pt}/\mathbb{C}^\times]$ and prove local cohomology theorems which imply they are well-defined.

In Section 6, we explain how to define Gromov-Witten invariants for $[X/\mathbb{C}^\times]$.

0.4. **Acknowledgements.** E.F. thanks Andrei Losev and Nikita Nekrasov for useful discussions. A.T. thanks Jarod Alper, Tom Coates, Dan Edidin, Ezra Getzler, Eduardo Gonzalez, Reimundo Heluani, and Chris Woodward for helpful conversations.

This research has been partially supported by DARPA and AFOSR through the grant FA9550-07-1-0543. In addition, E.F. and A.T. were supported by the NSF grant DMS-0303529, and C.T. was supported by the NSF grant DMS-0709448.

1. The Stack of Gieseker Bundles

In order to define the Gromov-Witten invariants of $[\mathrm{pt}/\mathbb{C}^\times]$ we need appropriate moduli stacks of marked curves carrying principal $\mathbb{C}^\times$-bundles. In this section, we introduce such stacks and discuss a few simple examples.

We always work over $\mathbb{C}$.

1.1. Curves.

In everything that follows, (C, σ_i) is a family of prestable marked curves, over a finitely-generated complex base scheme B. More precisely, $\pi : C \to B$ is a flat proper morphism whose fibers are connected complex projective curves of genus g with at worst nodal singularities, carrying a collection of marked points $\sigma_i : B \to C$ which are indexed by an ordered set I. A point σ_i is *special* if it is a node or a marked point. We shall always assume that any rational components of C have at least two special points.

We usually reserve the notation (Σ, σ_i) for families of *stable* marked curves. Recall that a marked curve is stable if each component of genus 0 carries at least 3 special points and each component of genus 1 carries at least 1 special point. $\overline{\mathcal{M}}_{g,I}$ denotes the stack of stable genus g, I-marked curves.

Notation 1.1. We will need to label three kinds of special points on C: ordinary marked points, nodes, and *trivialization points*. The latter are marked points at which we will trivialize the fibers of a $\mathbb{C}^\times$-bundle, but which are not required to be non-degenerate and do not count when determining stability. We will denote all such points by σ, distinguishing them by the subscript. Ordinary marked points are denoted σ_i, with $i \in I$. Nodes are σ_e, with e in a set E. Trivialization points are denoted σ_v, with v in a set V.

We use modular graphs to represent the topological type of a marked curve. Recall [BM96] that the *modular graph* γ of a curve (C, σ_i) consists of:

1. a vertex set V_γ (one vertex v for each component of C)
2. a half-edge set H_γ (one half-edge for each special point of C),
3. a gluing map $\partial_\gamma : H_\gamma \to V_\gamma$ (attaching half-edges to vertices),
4. an involution $j_\gamma : H_\gamma \to H_\gamma$, and
5. a function $g : V_\gamma \to \mathbb{N}$ (assigning to v the genus g_v of the normalization $\widetilde{C}_v$).

The involution j_γ generates an action of $\mathbb{Z}/(2)$ on H_γ, and the orbit set is a disjoint union of the set I_γ of tails (singlets, corresponding to marked points on C) and the set E_γ of edges (doublets, corresponding to nodes of C). The set E_γ may be further decomposed into the union

$$E_\gamma = E_\gamma^{split} \sqcup E_\gamma^{self}$$

of the set of splitting edges (which connect different vertices) and the set of self-edges (which start and end at the same vertex).



FIGURE 1. A (stable) marked curve and its associated modular graph, which has two splitting edges, one self-edge and one tail.

The substack of $\overline{\mathcal{M}}_{g,I}$ which classifies marked curves of type γ is denoted $\mathcal{M}_\gamma$. These substacks stratify $\overline{\mathcal{M}}_{g,I}$, in the sense that the closure of any $\mathcal{M}_\gamma$ is a union of modular graph-labeled substacks. The boundary divisors in this stratification have normal crossings.

1.2. **Bundles on Nodal Curves.** The quotient stack $[\mathrm{pt}/\mathbb{C}^\times]$ is, by definition, the classifying stack for principal $\mathbb{C}^\times$-bundles. Any principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$ determines and is determined by a map $\phi : C \to [\mathrm{pt}/\mathbb{C}^\times]$ such that the following diagram commutes.

$$\begin{tikzcd}
\mathcal{P} \arrow[r] \arrow[d, "p"] & \mathrm{pt} \arrow[d] \\
C \arrow[r, "\phi"] & {[\mathrm{pt}/\mathbb{C}^\times]}
\end{tikzcd}$$

The degree d of a principal $\mathbb{C}^\times$-bundle $\mathcal{P}$ is the degree $\phi_*[C] \in H_2^{\mathbb{C}^\times}(\mathrm{pt})$ of the associated morphism. The *multidegree* $\underline{d}$ of $\mathcal{P}$ assigns to each irreducible component $C_v \subset C$ the degree d_v of $\mathcal{P}|_{C_v}$.

$$\begin{aligned} \underline{d} : V_\gamma &\to H_2^{\mathbb{C}^\times}(\mathrm{pt}) \simeq \mathbb{Z} \\ v &\mapsto d_v, \end{aligned}$$

where V_γ is the set of vertices of the modular graph γ of C.

When reasoning about $\mathbb{C}^\times$-bundles on a fixed nodal curve C, we shall often use the following fact: Let $\nu : \widetilde{C} \to C$ denote the normalization of C. A principal bundle $\mathcal{P}$ on C determines and is determined by the following data:

1. a principal $\mathbb{C}^\times$-bundle $\widetilde{\mathcal{P}}$ $(= \nu^*\mathcal{P})$, and
2. for each node $\sigma \in \Sigma$, a *gluing isomorphism* $g : \widetilde{\mathcal{P}}_{\sigma^+} \simeq \widetilde{\mathcal{P}}_{\sigma^+}$, which identifies the fibers of $\widetilde{\mathcal{P}}$ over the preimages $\sigma^\pm$ of σ under ν.

Morally, the gluing isomorphiss are "transition functions" for two open sets whose intersection is the node σ.



FIGURE 2. Realizing $\mathcal{P}$ as $\widetilde{\mathcal{P}}$ together with a gluing datum g

This picture can also be used to understand the automorphisms of $\mathbb{C}^\times$-bundles. The automorphism group of the $\mathbb{C}^\times$-bundle $\widetilde{\mathcal{P}}$ is the product $(\mathbb{C}^\times)^V \times (\mathbb{C}^\times)^{2V'}$, where V is the set of stable components of C and V' is the set of rational components which have two special points. On a stable component of $\widetilde{C}$, we can rescale the fibers of $\widetilde{\mathcal{P}}$, while on each unstable rational component (assumed to have two special points), we can rescale the bundle and we can lift the $\mathbb{C}^\times$ of rotations of the curve. The automorphism group $(\mathbb{C}^\times)^V \times (\mathbb{C}^\times)^{V'}$ acts on the set of gluing isomorphisms, in the obvious way: For example, if we have a gluing isomorphism g which connects stable components $\widetilde{C}_v$ and $\widetilde{C}_{v'}$ (not necessarily different), then rescaling sends g to $g_v g g_{v'}^{-1}$.

An automorphism of $\widetilde{\mathcal{P}}$ gives rise to an automorphism of $\mathcal{P}$ if it fixes all the gluing isomorphisms. Note that, since $\mathbb{C}^\times$ is commutative, the gluing isomorphisms associated to self-nodes are automatically fixed by automorphisms of $\widetilde{\mathcal{P}}$.

1.3. **The Gieseker Completion.** By thinking about bundles on the normalization of a curve C, it is easy to see that $\mathbb{C}^\times$-bundles on nodal curves can become singular in families. The problem is that the space of gluing isomorphisms at a node $\sigma \in C$ is a copy of $\mathbb{C}^\times$; in families, these isomorphisms can approach the degenerate limits $\pm\infty$. More formally, the stack $\mathrm{Bun}_{\mathbb{C}^\times}(g, I)$ of $\mathbb{C}^\times$-bundles on stable marked curves of type (g, I) does not satisfy the valuative criterion for completeness.

We “fill in the holes” in $\mathrm{Bun}_{\mathbb{C}^\times}(g, I)$ by enlarging the classification problem slightly: We allow copies of $\mathbb{P}^1$ to appear at the nodes of stable curves, and insists that these $\mathbb{P}^1$ carry degree 1 bundles.

Definition 1.2. A morphism $m : C \to \Sigma$ of prestable curves is a *modification* if:

(1) m is an isomorphism on the complement of the preimage of the nodes of Σ, and
(2) the preimage under m of every node in Σ is either a node or a $\mathbb{P}^1$ with two special points.

If $\sigma \in \Sigma$ is a node and $m^{-1}(\sigma) \simeq \mathbb{P}^1$, we will say that Σ has been modified at σ.

A modification $m : C \to \Sigma$ is a *modification of I-marked curves* if $\sigma_i = m^{-1}(\sigma_i')$.

Note (a) that the notion of a modification of marked curves makes sense in families, (b) that modifications of marked curves do not introduce bubbles at marked points, only at nodes, and (c) that a smooth curve Σ has no non-trivial modifications.



Figure 3. A nodal curve Σ with a marked point σ_1 and its unique non-trivial modification C

Definition 1.3. A *Gieseker map to* $[\mathrm{pt}/\mathbb{C}^\times]$ is a pair $((C, \sigma_i), \mathcal{P})$ consisting of a prestable marked curve (C, σ_i) – whose stabilization we denote (Σ, σ_i) – and a principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$, for which

(1) the stabilization morphism $m : C \to \Sigma$ is a modification of I-marked curves, and
(2) For every node $\sigma \in \Sigma$ for which $m^{-1}(\sigma) \simeq \mathbb{P}^1$, the restriction $\mathcal{P}|_{m^{-1}(\sigma)}$ has degree 1.

The *degree* of a Gieseker bundle is the degree of the bundle $\mathcal{P}$.

We will frequently use the term “Gieseker bubble” for unstable $\mathbb{P}^1$’s carrying degree 1 bundles. Likewise, we will often say that the pair $(m : C \to \Sigma, \mathcal{P})$ is a “Gieseker bundle” on the stable curve (Σ, σ_i).

Example 1.4. Let (Σ, σ_i) be a stable curve of genus zero with four marked points. We assume Σ is nodal, with two components $\mathbb{P}^1_1$ and $\mathbb{P}^1_2$ meeting at an ordinary double point. (If Σ is smooth, Gieseker bundles are just ordinary bundles, which are classified by their degree and have automorphism group $\mathbb{C}^\times$.) Gieseker bundles on Σ come in two flavors:



Figure 4. Gieseker bundles on $\mathbb{P}^1 \cup \mathbb{P}^1$. Components are labelled with their degrees.

(1) Ordinary $\mathbb{C}^\times$-bundles on Σ: These are classified (up to isomorphism) by their multi-degrees $\underline{d} = (d_1, d_2)$. The automorphism group of any such bundle $\mathcal{P}$ is a copy of $\mathbb{C}^\times$, which rescales the fibers of $\mathcal{P}$.
(2) Gieseker bundles on the unique modification C of Σ: These are also classified by their multidegrees $\underline{d} = (d_1, 1, d_2)$. The automorphism group of any such bundle is a copy of $(\mathbb{C}^\times)^2$. One can rescale the bundle fibers, and one can lift the automorphisms of the unstable $\mathbb{P}^1$ to automorphisms of the bundle. (This is easily seen by working on the normalization $\widetilde{C}$.

Example 1.5. Let (Σ, σ_1) be the "boundary divisor" in $\overline{\mathcal{M}}_{1,1}$, a curve whose single rational component has one marked point (σ_1) and one self-node (σ).

Let's fix a total degree d. The degree d Gieseker bundles on (Σ, σ_1) come in two flavors, ordinary $\mathbb{C}^\times$-bundles on Σ and bundles on the modification of Σ at σ.



Figure 5. A pictures of (Σ, σ_1) and its modification (C, σ_1).

The normalization of Σ is a copy of $\mathbb{P}^1$. Line bundles on $\mathbb{P}^1$ are classified by their degree, so the ordinary degree d bundles on Σ are classified (up to isomorphism) by the gluing data associated to σ. The space of such gluing data is a copy of $\mathbb{C}$. σ is a self-node, so these gluing isomorphisms are fixed by bundle rescalings. Hence the automorphism group of an ordinary bundle on Σ is a copy of $\mathbb{C}^\times$.

The modification C has two rational components and two nodes, so a bundle on C is specified by two gluing isomorphisms. However, these gluing isomorphism do not classify; we can rescale the bundle differently on each component of the normalization $\widetilde{C}$ and we we can lift the rotations of the Gieseker bubble. A rescaling and the rotations can be used to fix the gluing isomorphisms, so (up to isomorphism) there is only *one* bundle on the modification which satifies the Gieseker conditions. The automorphism group of this bundle is a copy of $\mathbb{C}^\times$, coming from the remaining automorphism of the bundle data on the normalization.

Remark 1.6. The following story, though not rigorous, can help to understand the intuition behind these definitions.

Fix a nodal curve Σ and suppose that we have a family of $\mathbb{C}^\times$-bundles on Σ, parametrized by a coordinate t, for which the gluing isomorphism $g : \mathcal{P}_{\sigma^+} \to \mathcal{P}_{\sigma^-}$ over a node $\sigma \in \Sigma$ approaches 0 at $t \to 0$. (Assuming that $g \to 0$ is no loss of generality; the other limit $g \to \infty$ is equivalent to $g^{-1} \to 0$.) We want to replace this singular limit with a bundle defined on some other curve C.

One can guess how to do this by looking at a section s of an associated fiber bundle $V = \mathcal{P} \times_{\mathbb{C}^\times} \mathbb{C}$. If we lift s to a section $\tilde{s}$ on the normalization, it must obey

$$\tilde{s}(\sigma^+) = g\tilde{s}(\sigma^-).$$

In this limit, we must have $\tilde{s}(\sigma^+) \to 0$. By continuity, the section s on Σ must have a zero which approaches the node as $g \to 0$. (When $g \to \infty$, the zero approaches the node from the other side.)

To keep track of how a single zero z approaches the node, we should replace the singular limit with a new bundle on C obtained by creating a $\mathbb{P}^1$ at the node. (This $\mathbb{P}^1$ records the way the zero approached the node.) The section on this new component must have one zero and no poles, so the degree of the new bundle on this component must be 1. The total degree of the bundle is a topological invariant, so the degree of the bundle on the original component will drop by 1.



FIGURE 6. A zero approaches at node, leading to a Gieseker bubble

Definition 1.7. $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is the fibered category (over $\mathbb{C}$-schemes) which classifies Gieseker bundles on stable genus g, I-marked curves. Its objects are tuplets $(B, C, \sigma_i, \mathcal{P})$ consisting of

- a test scheme B,
- a flat projective family $\pi : C \to B$ of pre-stable I-marked, genus g curves with marked points $\sigma_i : B \to C$, and
- a principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$

such that for every closed point $b \in B$, the pair $((C_b, \sigma_{b,i}), \mathcal{P}|_{C_b})$ is a Gieseker map to $[\mathrm{pt}/\mathbb{C}^\times]$. The morphisms in this category are Cartesian diagrams

$$\begin{tikzcd}
\mathcal{P}' \arrow[r, "\tilde{f}"] \arrow[d, "p'"] & \mathcal{P} \arrow[d, "p"] \\
C' \arrow[r, "f"] \arrow[d, "\pi'"] & C \arrow[d, "\pi"] \\
B' \arrow[r] & B
\end{tikzcd},$$

where $\tilde{f}$ is $\mathbb{C}^\times$-equivariant and $\sigma_i = f \circ \sigma'_i$.

Notation 1.8. The *degeneration type* of a pair $((C, \sigma_i), \mathcal{P})$ is the pair $(\gamma, \underline{d})$ consisting of the modular graph γ of (C, σ_i) and the multidegree $\underline{d}$ of $\mathcal{P}$. $\mathcal{M}_{\gamma,\underline{d}}$ is the substack of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ which classifies bundles of degeneracy type $(\gamma, \underline{d})$.

Remark 1.9. The condition that a pair $((C, \sigma_i), \mathcal{P})$ be a Gieseker map can be phrased purely in terms of the degeneracy type.

In Section 2, we will introduce an atlas for $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ (thereby proving it to be a stack) and discuss its geometric properties. In the remainder of this section, we introduce notation for our stack's universal families and its modular graph decomposition, and then discuss a few key examples.

1.4. **Universal Families.**

By virtue of its definition, $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ carries several universal families. It carries a family of semi-stable curves

$$\pi : C \to \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$$

with marked points

$$\sigma_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to C$$

indexed by I, and it has a principal $\mathbb{C}^\times$-bundle

$$p : \mathcal{P} \to C.$$

The universal $\mathbb{C}^\times$-bundle defines a universal map

$$\phi : C \to [\mathrm{pt}/\mathbb{C}^\times]$$

and the compositions $\mathrm{ev}_i = \phi \circ \sigma_i$ are the *evaluation maps*

$$\mathrm{ev}_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to [\mathrm{pt}/\mathbb{C}^\times] \qquad (i \in I).$$

Finally, there is a natural morphism

$$F : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to \overline{\mathcal{M}}_{g,I}$$

which forgets the principal bundle $\mathcal{P}$ and sends the semi-stable marked curve (C, σ_i) to its stabilization.

1.5. **Examples.** We treat three examples here. The first is trivial. The second is the simplest example in which the Gieseker bubbles can be seen. The third illustrates how the substacks classifying non-trivial Gieseker bundles "fill in the gaps" between the substacks which parametrize bundles which live on the same nodal curve but have different multidegrees.

Example 1.10 (genus zero, 3 marked points)**.** $\overline{\mathcal{M}}_{0,3}$ is a point; there is, up to equivalence, only one stable genus zero curve with 3 marked points. On this curve, up to equivalence, there is one bundle of degree d; any $d \in \mathbb{Z}$ is allowed. The automorphism group of $\mathcal{O}_{\mathbb{P}^1}(d)$ is a copy of $\mathbb{C}^\times$, so we conclude

$$\widetilde{\mathcal{M}}_{0,3}([\mathrm{pt}/\mathbb{C}^\times]) \simeq \bigsqcup_{d\in\mathbb{Z}}[\mathrm{pt}/\mathbb{C}^\times].$$

Example 1.11 (genus one, 1 marked point)**.** This example shows the Gieseker completion at work.

Let's fix the total degree d of the Gieseker bundles. We'll represent the stack $\widetilde{\mathcal{M}}^d_{1,1}([\mathrm{pt}/\mathbb{C}^\times])$ of degree d Gieseker bundles as a quotient $[A/\mathbb{C}^\times]$, where A is the stack which classifies pairs $((C,\mathcal{P}),t)$ consisting of a Gieseker bundle and a trivialization $t : \mathcal{P}_\sigma \simeq \mathbb{C}^\times$ of the Gieseker bundle at the marked point σ. A also comes equipped with a forgetful map $f : A \to \overline{\mathcal{M}}_{1,1}$.

Over smooth curves (Σ, σ_1), the fibers of f are copies of the Jacobian $\mathrm{Jac}(\Sigma)$ of Σ. The Jacobian of Σ is a copy of Σ, so over $\mathcal{M}_{1,1}$, the stack A is simply a copy of the universal curve $\Sigma_{1,1}$. In fact, this is also true over the boundary divisor of $\overline{\mathcal{M}}_{1,1}$. We saw in Example 1.5 that the space of Gieseker bundles (with a trivialization t to eliminate the global rescaling automorphisms) is obtained by gluing a copy of the point pt to a copy of $\mathbb{C}^\times$. This gluing identifies the pt with both 0 and ∞, so the resulting curve is a copy of (Σ, σ_1).

Example 1.12 (genus zero, 4-marked points)**.** Recall that $\overline{\mathcal{M}}_{0,4}$ is isomorphic to $\mathbb{P}^1$. The open locus classifying smooth curvces is $\mathbb{P}^1 \setminus \{0, 1, \infty\}$, and the boundary divisor $\{0, 1, \infty\}$ classifies reducible nodal curves with two marked points on each

Let $B = \mathbb{P}^1 \setminus \{1, \infty\} = \mathbb{A}^1 \setminus 1$ and consider the family $(\Sigma, \sigma_i) : B \hookrightarrow \overline{\mathcal{M}}_{0,4}$ of marked curves obtained by restricting the universal marked curve $\Sigma_{0,4}$ to B. This family is a deformation of the curve $(\Sigma_o, \sigma_{o,i})$ which has two components meeting at a common node and each carrying two marked points.

We will describe the stack $\widetilde{\mathcal{M}}_{0,4}([\mathrm{pt}/\mathbb{C}^\times])$ by giving an atlas A for the restriction $F|_B : \widetilde{\mathcal{M}} \to B$ of $F : \widetilde{\mathcal{M}}_{0,4}([\mathrm{pt}/\mathbb{C}^\times]) \to \overline{\mathcal{M}}_{0,4}$ to B. (Nearly identical descriptions apply near 1 and ∞ in $\overline{\mathcal{M}}_{0,4}$, and $\widetilde{\mathcal{M}}_{0,4}([\mathrm{pt}/\mathbb{C}^\times])$ is obtained by gluing these local descriptions.) Let V be the subset $\{0, \infty\} \subset I$. The atlas A is the algebraic space (scheme, in fact) which classifies isomorphism classes of tuplets

$$(C, \sigma_i, \mathcal{P}, t_0, t_\infty)$$



where $(C, \sigma_i, \mathcal{P})$ is a family of Gieseker bundles on the curve Σ/B and the $t_v \in \Gamma(B, \sigma_v^*\mathcal{P})$ are families of trivializations based at the marked points 0 and ∞. (For any closed point $b \in B$, $t_v(b)$ is a point in the fiber of $\mathcal{P}$ at $\sigma_v(b)$.)

A is a disjoint union

$$A = \sqcup_{d\in\mathbb{Z}} A_d$$

of isomorphic subschemes which classify Gieseker bundles of total degree d. In the remainder of this example, we fix a total degree d, and focus on A_d.

Gieseker bundles on the fibers of $\pi : \Sigma \to B$ were classified in Example 1.4. Adding the two trivializations eliminates the automorphisms, and introduces an extra degree of freedom on the unmodified curves, which may be thought of as the ratio of the two trivializations. Thus, the generic fiber of the forgetful morphism $f : A_d \to B$ is isomorphic to $\mathbb{C}^\times$, while the special fiber is an endless chain of rational curves $\cup_{n\in\mathbb{Z}}\mathbb{P}^1_n$. The open $\mathbb{C}^\times_n \subset \mathbb{P}^1_n$ classifies bundles-with-trivializations whose multidegree is $(d + n, -n)$. The non-empty intersection points $\mathrm{pt}_n = \mathbb{P}^1_{n-1} \cap \mathbb{P}^1_n$ classify bundles-with-trivializations whose multidegree is $(d+n-1, 1, -n)$. (For reasons of space, we have drawn only finitely many of the $\mathbb{P}^1$'s in the figure below.)



The action of the group $(\mathbb{C}^\times)^V = \mathbb{C}^\times_0 \times \mathbb{C}^\times_\infty$ which rescales the trivializations preserves the forgetful morphism f. The diagonal $\mathbb{C}^\times_\Delta \subset (\mathbb{C}^\times)^V$ acts trivially.

For later computations, it is useful to have an explicit Cech cover of A. Let

$$A_{d,n} = \operatorname{Spec}\mathbb{C}[z_n, w_n] \simeq \mathbb{A}^2.$$

We obtain A_d by gluing $A_{d,n-1}$ to $A_{d,n}$, identifying the open sets U_{w_n} and $U_{z_{n-1}}$ via the relation $w_n = 1/z_{n-1}$. (Here $U_{w_n} = \operatorname{Spec} \mathbb{C}[z_n, w_n]_{(w_n)} \simeq \mathbb{A} \times \mathbb{C}^\times$; likewise, $U_{z_{n-1}}$.) The morphism $f : A \to B$ is given by $b = z_n w_n$, where b is the standard coordinate on $B \subset \operatorname{Spec} \mathbb{C}[b]$. In this notation, pt_n is the origin, $\mathbb{C}_n^\times$ is the z_n coordinate axis, and $\mathbb{C}_{n-1}^\times$ is the w_n coordinate axis. Finally, $(\mathbb{C}^\times)^V \simeq (\mathbb{C}^\times)^2$ acts on A_d, with weight $(1, -1)$ on z_n and weight $(-1, 1)$ on w_n. Thus, the diagonal acts trivially, and the fixed points are picked out by $z_n = w_n = 0$. One may think that $z_n \propto w_n^{-1} \propto t_0/t_\infty$, at least generically.

2. The Geometry of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$

In this section, we describe the geometry of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$.

We exhibit an atlas for our stack. We will use this atlas – denoted A – in the proof that the Gromov-Witten invariants of $[\mathrm{pt}/\mathbb{C}^\times]$ are well-defined. While discussing atlases, we also prove that the stack of Gieseker bundles satisfies a valuative criterion for completeness.

Second, we discuss the deformations of Gieseker bundles. We use this information to describe the connected components of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ and to give $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ a stratification by degeneracy type.

2.1. **Our Favorite Atlas.**

The atlas defined in Proposition 2.2 below generalizes the ones in Examples 1.11 and 1.12.

Let $(\Sigma_o, \sigma_{o,i})$ be a point in $\overline{\mathcal{M}}_{g,I}$, with modular graph γ_o, and let B be an affine étale coordinate chart centered at $(\Sigma_o, \sigma_{o,i})$, represented by a curve (Σ, σ_i) over B. We can assume with no loss of generality that the modular graph strata of B coincide with the coordinate axes, so that $(\Sigma_o, \sigma_{o,i})$ is the most degenerate curve in the family. $\overline{\mathcal{M}}_{g,I}$ is covered by charts of this form.

We denote the fiber of F over B by $\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$. This stack classifies Gieseker bundles on the family (Σ, σ_i).

$$\begin{tikzcd}
\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)} \arrow[r] \arrow[d] & \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \arrow[d, "F"] \\
B \arrow[r] & \overline{\mathcal{M}}_{g,I}
\end{tikzcd}$$

We want to exhibit an atlas for $\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$. In the moduli theory of bundles, one often obtains atlases by fixing one or more marked points and then parametrizing bundles equipped with trivializations at these marked points. We would like to do this in families.

Definition 2.1. Suppose that C comes equipped with a marked point $\sigma : B \to C$. A *family of trivializations based at* σ is a section $t \in \Gamma(B, \sigma^*\mathcal{P})$.

Such families of marked points induce isomorphisms $t_{\sigma(b)} : \mathcal{P}_{\sigma(b)} \simeq \mathbb{C}^\times$ for every geometric point $b \in B$.

Proposition 2.2. (1) *Let $V = V_{\gamma_o}$ be the vertex set of the modular graph of $(\Sigma_o, \sigma_{o,i})$. After étale refinement of the neighborhood B, we may choose a collection of marked points $\sigma_v : B \to \Sigma$ (with $v \in V$) such that every component of every geometric fiber of Σ carries at least one of the σ_v.*

(2) *Let $A = A_{(\Sigma,\sigma_i)}(\{\sigma_v\})$ denote the stack which classifies triplets*

$$((C, \sigma_i), p : \mathcal{P} \to C, t_v \in \Gamma(B, \sigma_v^*\mathcal{P})\}_{v \in V_{\gamma_o}})$$

of Gieseker bundles with trivializations at the selected marked points. A is an algebraic space. Moreover the natural action of $\mathcal{G} = (\mathbb{C}^\times)^V$, with the v-th copy $\mathbb{C}^\times_v$ rescaling the v-th trivialization, displays $\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$ as the quotient stack

$$\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)} = [A/\mathcal{G}].$$

Proof. The first claim follows from the local geometry of the stack of marked curves. If we choose one new marked point σ_v on each component of Σ_o, we obtain a point in the moduli stack $\overline{\mathcal{M}}_{g,I\sqcup V}$. After étale refinement $B' \to B$, there are no obstructions to lifting the morphism $B' \to \overline{\mathcal{M}}_{g,I}$ to a morphism $B' \to \overline{\mathcal{M}}_{g,I\sqcup V}$.

To prove the second claim, we observe that the $(\mathbb{C}^\times)^V$ action preserves the fibers of the forgetful map $f : A \to B$. Representing a bundle on C as a bundle on $\widetilde{C}$ together with gluing isomorphisms, it is easy to see that the trivializations eliminate all of the automorphisms of the Gieseker bundles. It follows that A is an algebraic space. The action of $(\mathbb{C}^\times)^V$ is freely transitive on the trivializations, so quotienting by this action is equivalent to forgetting the trivialiations. □

Corollary 2.3. *$\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is a smooth Artin stack.*

Proof. Choose a collection $\{B_\alpha\}$ of étale neighborhoods which cover $\overline{\mathcal{M}}_{g,I}$. The collection $\{A_\alpha\}$ covers $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$. □

Corollary 2.4. *The dimension of $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is*

$$dim(\mathbb{C}^\times)(g-1) + 3(g-1) + |I|.$$

Proposition 2.5. *The stack $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ satisfies the "valuative criterion for completeness": Let R be a complete discrete valuation ring, with fraction field K, and consider the "disc" $D = \mathrm{Spec}(R)$ and the "punctured disc" $D^\times = \mathrm{Spec}(K)$. Given a family $((C, \sigma_i), \mathcal{P})$ of Gieseker bundles on $D^\times$, we can extend the family to $D^\times$ (possibly after étale base change).*

Proof. Although rarely stated in this form, this result is implicit in the literature on GIT constructions of coarse moduli spaces. To keep this paper somewhat self-contained, we sketch a proof.

$\overline{\mathcal{M}}_{g,I}$ is complete, so it's enough to prove the valuative criterion for the forgetful morphism $F : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}\,/\mathbb{C}^\times]) \to \overline{\mathcal{M}}_{g,I}$. We fix a stable curve (Σ, σ_i) extending the stabilization of C to (an étale refinement of) D.

The result now follows from the projectivity of coarse moduli spaces of S-equivalence classes of families of rank 1 torsion free sheaves on varying nodal curves. More precisely: A result of Seshadri [Ses82] implies that there is a surjective morphism from the fiber of F over the chosen family $(\Sigma, \sigma_i) : D \to \overline{\mathcal{M}}_{g,I}$ onto the stack of stable rank 1 torsion free sheaves on the family Σ; we associate to any principal $\mathbb{C}^\times$-bundle $\mathcal{P}$, the pushforward m_*E of the line bundle E associated to $\mathcal{P}$ and the standard representation of $\mathbb{C}^\times$. Simpson's GIT construction [Sim94], based on Maruyama's boundedness results [Mar81], guarantees that the coarse algebraic moduli space of S-equivalence classes of rank 1 torsion-free sheaves on a family of projective curves is projective and hence complete. Completeness of the coarse moduli space implies the existence (though not uniqueness) of some family of torsion-free sheaves extending any m_*E, hence the existence (though not uniqueness) of a family of Gieseker bundles extending the given one. □

The stack $\overline{\mathcal{M}}_{g,I}([\mathrm{pt}\,/\mathbb{C}^\times])$ is not separated, because the degree d of a bundle can split as $d = (d+n) - n$ for any $n \in \mathbb{Z}$ if the curve develops a node. However, this is essentially the only source of non-separatedness.

Proposition 2.6. *The fiber of F at any nodal curve (Σ, σ_i) in $\overline{\mathcal{M}}_{g,I}(\mathrm{pt})$ is separated.*

2.2. **Deformations.**

First, we want to show the existence of deformations.

Proposition 2.7. *$\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}\,/\mathbb{C}^\times])$ is unobstructed.*

Proof. The obstructions to deforming a bundle $\mathcal{P}$ on a fixed curve C are captured by the Ext-group $\mathrm{Ext}^2((p_*L_{\mathcal{P}})^{\mathbb{C}^\times}, \mathcal{O}_C)$. Here $\mathcal{O}_C$ is the structure sheaf of C and $(p_*L_{\mathcal{P}})^{\mathbb{C}^\times}$ is the $\mathbb{C}^\times$-invariants in the pushdown of the cotantent complex of $\mathcal{P}$ along the structure morphism $p : \mathcal{P} \to C$. This Ext-group vanishes because C is one-dimensional. The vanishing of obstructions to deforming the curve and the bundle together follows from the tangent-obstruction sequence derived from the exact triangle associated to the cotangent complex $L_{\mathcal{P}/C}$. □

It follows that any Gieseker bundle $(C_o, \sigma_{o,i}, \mathcal{P}_o)$ can be deformed to a family $(C, \sigma_i, \mathcal{P})$ on the Spec of a discrete valuation ring.

The following lemma characterizes the degeneracy types which can occur when we deform a given Gieseker bundle.

Lemma 2.8. *Let $(C_o, \sigma_{o,i}, \mathcal{P}_o)$ be a Gieseker bundle having degeneracy type $(\gamma_o, \underline{d}_o)$, and suppose that we are given a deformation $(C, \sigma_i, \mathcal{P})$ of $(C_o, \sigma_{o,i}, \mathcal{P}_o)$ over the Spec of a discrete valuation ring. The degeneracy type $(\gamma, \underline{d})$ of the generic fiber can be any degree-labelled modular graph obtained from $(\gamma_o, \underline{d}_o)$ by the combinations of the following elementary operations:*

(1) *Resolve a self node: Delete a self-edge attached to a vertex* v*, increasing the genus* g_v *by* 1*, leave the multi-degree unchanged.*
(2) *Resolve a splitting node: replace a pair of vertices* v_1 *and* v_2 *with a single vertex* v*, having genus* $g_v = g_{v_1} + g_{v_2}$ *and multidegree* $d_v = d_{v_1} + d_{v_2}$*. Any other splitting edges connecting* v_1 *and* v_2 *become self-edges.*

Moreover all such modular graphs occur in some deformation.

Proof. Induction on the number of nodes, using the modular graph stratification of the stack of curves. □

Corollary 2.9. $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ *is locally of finite type.*

Proof. Every stratum is clearly of finite type, and the lemma says that we only reach finitely many strata by deformation. □

Corollary 2.10. *The connected components of* $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ *are labelled by total degree* d *of the Gieseker bundles.*

$$\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) = \bigsqcup_{d\in\mathbb{Z}} \widetilde{\mathcal{M}}^d_{g,I}([\mathrm{pt}/\mathbb{C}^\times]).$$

Remark 2.11. The corollary makes it obvious that $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is of infinite type. The connected components $\widetilde{\mathcal{M}}^d_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ can also be of infinite type. Any modular graph γ with at least two vertices carries countably many multi-degrees $\underline{d} : V_\gamma \to \mathbb{Z}$ for which $\sum_{v\in V_\gamma} d_v = d$.

Corollary 2.12. *The modular graph decomposition (see Definition 1.8)*

$$\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) = \bigcup_{(\gamma,\underline{d})} \mathcal{M}_{(\gamma,\underline{d})},$$

is a stratification, meaning that the closure of $\mathcal{M}_{(\gamma,\underline{d})}$ *in* $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ *is obtained as a union*

$$\mathrm{cl}\left(\mathcal{M}_{(\gamma,\underline{d})}\right) = \bigcup_{(\gamma',\underline{d}')} \mathcal{M}_{(\gamma',\underline{d}')},$$

where the primed union is over all multidegree-labelled modular graphs $(\gamma,\underline{d}')$ *obtained from* $(\gamma,\underline{d})$ *by sequences of the following elementary operations:*

(1) *Self node: Lower the genus of a vertex by 1, and add a self-edge.*
(2) *Splitting node: Split a vertex* v *into two vertices* v_1 *and* v_2*, connected by an edge, with* $g_{v_1} + g_{v_2} = g_v$ *and* $d_{v_1} + d_{v_2} = d_v$*.*
(3) *Gieseker bubbling: Replace an edge connecting a stable vertex* v *to a stable vertex* v' *with two edges connected to a common vertex* v *having* $g_v = 0$ *and* $d_v = 1$*, while subtracting* 1 *from the degree* d_v *or* $d_{v'}$*. (Note that* v *may equal* v'*.)*

Proposition 2.13. *The boundary of the closure in the corollary above is a divisor with normal crossings.*

3. The Structure of the Local Atlas

In this section, we describe some aspects of the geometry of our atlas A which we will need in the proof of our main theorem.

We begin by introducing a refinement of A's modular graph stratification. Then we study the action of $\mathcal{G} = (\mathbb{C}^\times)^V$ on A, and characterize the stabilizer groups of the points of A. In particular, we shall see that the Kirwan-Ness stratification by stabilizer group is compatible with our refinement of the modular graph stratificaiton.

Next, we identify some finite-type subspaces S_{N_u,N_l} of A, which parametrize Gieseker bundles on the family Σ/B whose multidegrees obey certain bounds involving collections N_u and N_l of integers. We show that changing N_u or N_l amounts to adding or deleting strata which parametrize certain deformations of the fixed point loci of subgroups of $\mathcal{G}$.

Finally, we show that, if N_u and N_l are "as close as possible", then the quotient stack

$$[S_{N_u,N_l}/(\mathbb{C}^\times)^V] \simeq Q_N \times [\mathrm{pt}/\mathbb{C}^\times_\Delta],$$

where $\mathbb{C}^\times_\Delta \subset (\mathbb{C}^\times)^V$ is the diagonal subgroup and Q_N is an algebraic space which is proper over B.

Notation 3.1. In the remainder of this section, A is the component of the atlas A_α associated to a particular étale neighborhood B in $\overline{\mathcal{M}}_{g,I}$, centered at a curve whose modular graph is γ_o.

3.1. **Refined Modular Graphs.**

When discussing the deformations of a curve, it can be useful to keep track of which nodes of the curve are being smoothed. We introduce the following definition to make this idea precise.

Definition 3.2. A *deformation γ of the modular graph γ_o* consists of the following data:

(1) a subset $E_\gamma \subset E_{\gamma_o}$, and
(2) a partition V_γ of $R_{\gamma_o} = \sqcup_{v \in V_\gamma} V^v_{\gamma_o}$.

These data determine a modular graph (also called γ), whose vertices are the blocks of the partition V_γ, with edge set E_γ and tail set I_{γ_o}. The gluing maps come from γ_o. The set of splitting edges E^{split}_γ of a deformation of a modular graph is the set of edges in E_γ which connect different blocks of the partition, i.e., the splitting edges of the modular graph γ.

A deformation of a multi-degree labelled modular graph $(\gamma_o, \underline{d}_o)$ is simply a deformation of the underlying modular graph, together with a labelling $\underline{d} : V_\gamma \to \mathbb{Z}$ satisfying the conditions enumerated in Proposition 2.8.

Proposition 3.3. *The stratification of A by labelled modular graphs can be refined to a stratification by deformations of labelled modular graphs (possibly after restricting to an smaller étale neighborhood $B' \to B$).*

3.2. **Stabilizer Groups.** Let $(C, \sigma_i, \mathcal{P}, \{t_v\})$ be a Gieseker bundle representing a closed point of A, i.e., (C, σ_i) is a modification of the fiber of $\pi : \Sigma \to B$ over some closed point $b \in B$. We say that a group element $(g_v) \in (\mathbb{C}^\times)^V$ *stabilizes* $(C, \mathcal{P}, \{t_v\})$ if the action of (g_v) – given by $t_v \mapsto g_v t_v$ – produces an isomorphic object.

We identify the stabilizer group of $(C, \sigma_i, \mathcal{P}, \{t_v\})$ by working on the normalization $\widetilde{C}$, as explained in Section 1.2; the bundle $\mathcal{P}$ is specified in terms of a bundle $\widetilde{\mathcal{P}}$ and a collection of gluing isomorphisms $\{g_e\}$, one for each node σ_e in C. The trivializations simply lift; each stable component of $\widetilde{C}$ carries one or more trivializations.

We can use this picture to identify the automorphisms of $(C, \sigma_i, \mathcal{P}, \{t_v\})$ as follows: If we rescale a trivialization t_v by the group element $g_v \in \mathbb{C}^\times_v$, then we can try to compensate this rescaling with an automorphism of the bundle $\widetilde{\mathcal{P}}$. There are two obstacles to carrying out this procedure:

1. rescaling $\widetilde{\mathcal{P}}$ on $\widetilde{C}_v$ may rescale other trivializations, and
2. rescaling may change the gluing maps, so that the automorphism of $\widetilde{\mathcal{P}}$ doesn't descend to an automorphism of $\mathcal{P}$.

We can only avoid the first obstacle if we act with the same group element on any trivializations which are based on the same component of $\widetilde{C}$. The second obstacle can be avoided in two ways: If a gluing isomorphism connects the stable component $\widetilde{C}_v$ to a Gieseker bubble, we can use the extra automorphism of the Gieseker (lifting the bubble's rotations) to rescale this gluing isomorphism, while leaving the gluing isomorphism on the other side of the bubble untouched. Alternately, if the gluing isomorphism connects two stable components, we must act on all trivializations on these components with the same group element.

We formalize this discussion as follows:

Definition 3.4. Let $\gamma = (E_\gamma, V_\gamma)$ be a deformation of the modular graph γ_o. The *partition $R = R(\gamma)$ of $V = V_{\gamma_o}$ associated to* γ is the one given by the equivalence relation which identifies v and v' if there is a path in γ from the vertex represented by the block $\partial_{\gamma_o}(v)$ to the vertex represented by the block $\partial_{\gamma_o}(v')$, which does not pass through any unstable node.

Definition 3.5. Let R be a partition of the $V = V_{\gamma_o} = \sqcup_{r \in R} V^r$. $(\mathbb{C}^\times)^R$ denotes the subgroup $(\mathbb{C}^\times)^{V^r} \subset (\mathbb{C}^\times)^V$ corresponding to the block V^r, and $\mathbb{C}^\times_{\Delta^r} \subset (\mathbb{C}^\times)^{V^r}$ denotes the diagonal subgroup of $(\mathbb{C}^\times)^{V^r}$. The *subgroup $\mathcal{G}_R \subset \mathcal{G}$ associated to the partition R* is the product

$$\mathcal{G}_R = \Pi_{r \in R} \mathbb{C}^\times_{\Delta^r}.$$

Proposition 3.6. *Let $(C, \sigma_i, \mathcal{P}, \{t_v\})$ be a closed point of the stratum $A_{\gamma,\underline{d}}$ labelled by a deformation $\gamma = (E_\gamma, V_\gamma)$ of γ_o and a multi-degree $\underline{d}$. The stabilizer group of $(C, \sigma_i, \mathcal{P}, \{t_v\})$ is the group $\mathcal{G}_R$ associated to the partition $R = R(\gamma)$.*

Note that the stabilizer group depends only on the modification γ and not on the multi-degree.

Note also that $\mathcal{G}_R$ always contains the diagonal subgroup $\mathbb{C}^\times_\Delta$ of $(\mathbb{C}^\times)^V$; this subgroup is the group of global rescalings of $\mathcal{P}$.

Definition 3.7. We say that a partition is *non-trivial* if $\mathcal{G}_R \neq \mathbb{C}^\times_\Delta$. We denote the collection of edges which link different blocks of the partition by E_R^{split}.

3.3. **The Fixed Point Loci.**

Let $F_R \subset A$ denote the fixed point locus of the group $\mathcal{G}_R$, i.e., the substack of A classifying Gieseker bundles whose stabilizer group is $\mathcal{G}_R$. The stabilizer group of a Gieseker bundle depends only on the refined modular graph, so F_R is a union of strata $A_{\gamma,\underline{d}}$. Thus, the stratification of A by stabilizer groups is compatible with the stratification by deformations of labelled modular graphs.

Deformations which remain in the fixed point locus F_R must not smooth any of the Gieseker bubbles which define the partition R. This implies that such deformations must preserve the partial sums

$$n_r = \sum_{v \in V^r_{\gamma_o}} d_v, \qquad r \in R.$$

It is easy to see that the connected components of F_R are in correspondence with collections of integers $\underline{n} = (n_r)_{r\in R} \in \mathbb{Z}^R$.

$$F_R = \bigsqcup_{\underline{n}} F_R^{\underline{n}}.$$

Roughly speaking, these integers $\underline{n}$ keep track of the multidegrees on the stable components of C.

Example 3.8. In the special case of Example 1.12, the only non-trivial partition is the two-block partition $R = \{\{v_+\},\{v_-\}\}$ associated to the modular graph of the special fiber $(\Sigma_o, \sigma_{o,i})$. The fixed point stratum $F_R^{\underline{n}} \in A_d$ labelled by $\underline{n} = (n_+, n_-)$ is the point $\mathrm{pt}_{\underline{n}}$ which classifies Gieseker bundles-with-trivializations whose multidegree is $(n_+, 1, n_-)$. (In Example 1.12, we used the notation $\mathrm{pt}_n \in A_d$, with $d = n_+ + n_- + 1$ and $n = -n_-$.)

3.4. **Finite-Type Subspaces of A.**

It follows from the discussion in Section 1 that the connected components A_d of the atlas A are of infinite type whenever $|V_{\gamma_o}| \geq 2$, and finite type otherwise. When $|V_{\gamma_o}| = 1$, the stack A_d is proper, relative to B; it is the compactified Jacobian of Σ. (Indeed, it is clearly of finite type and complete. And, because Σ/B is a family of one-component curves, there is no degree-splitting, so A^d is separated in this case.)

However, when $|V_{\gamma_o}| \geq 2$, the atlas A_d of has infinitely many finite-type strata. It is useful to think of this proliferation of strata as follows: The fiber of $f : A \to B$ over a point $b \in B$ for which Σ_b has only self-nodes is of finite-type. But if b

moves in such a way that Σ_b splits into two components, then the degree d can split between these components arbitrarily, as $d = (d+n) - n$. This can happen in multiple ways, depending on how Σ_b acquires nodes; moreover, the corresponding strata can meet in higher codimension, if Σ_b degenerates to a curve having multiple components.

In this section, we define some finite-type algebraic subspaces $S_{N_u,N_l} \subset A_d$. These stacks parametrize Gieseker bundles whose multidegrees obey certain inequalities, which govern how d can split as $(d+n)-n)$.

Notation 3.9. When R is a non-trivial two-block partition of V_{γ_o} (with blocks V^+ and V^-), the fixed point strata $F_R^{\underline{n}}$ are labelled by pairs of integers $\underline{n} = (n_+, n_-)$. On A_d, the total degree d is fixed and we must have $n_+ + n_- + k = d$, where $k = k(R) = |E_R^{split}|$ is the number of edges in γ_o which connect a vertex in V^+ to a vertex in V^-. In this situation, it makes sense to label fixed point strata with a single integer N, for which $\underline{n} = (n_+, n_-) = (d+N-k, -N)$. Making this choice simplifies some of our notation, and introduces funny asymmetries elsewhere.

Definition 3.10. Choose collections of integers

$$N_u = (N_u(R)) \qquad \text{and} \qquad \underline{N}_l = (N_l(R))$$

where $R = \{V_{\gamma_o}^+, V_{\gamma_o}^-\}$ ranges over the set $NT2B(\gamma_o)$ of non-trivial 2-block partitions of $V = V_{\gamma_o}$. We assume that $N_u(R) > N_l(R)$ for all R.

If $(\gamma, \underline{d})$ is any multidegree-labelled deformation of γ_o which has at least two vertices, then the choice of a 2-block partition R of V_{γ_o} induces a 2-block partition of $V_\gamma = V_\gamma^+ \sqcup V_\gamma^-$. We define, for any 2-block partition R, the partial sums

$$d_+(\gamma, R) = \sum_{v \in V_\gamma^+} d_v \qquad d_-(\gamma, R) = \sum_{v \in V_\gamma^-} d_v.$$

The *locus* S_{N_u,N_l} *of degree* d *Gieseker bundles with multidegree bounded by* N_u *and* N_l is the union of all strata $A_{\gamma,\underline{d}}$ in A_d, for which, if γ has multiple vertices, then for any partition $R \in NT2B(\gamma_o)$, the partials sums defined above obey the following inequalities:

$$\begin{aligned} d_+(\gamma, R) \geq & d + N_l(R) - k(R) + 1 \\ d_-(\gamma, R) \geq & - N_u(R) + 1, \end{aligned}$$

where $k(R) = |E_R^{split}|$ is the number of edges in γ_o connecting vertices in $V_{\gamma_o}^+$ to vertices in $V_{\gamma_o}^-$.

Note that

(1) If $N_u'(R) \geq N_u(R)$ and $N_l'(R) \leq N_l(R)$, then

$$S_{N_u,N_l} \subset S_{N_u',N_l'}.$$

(2) A_d is the increasing limit

$$A_d = \lim_{N_u(R)\to\infty, N_l(R)\to -\infty} S_{N_u,N_l}$$

Example 3.11. In the special case of Example 1.12, the atlas A_d is finite-type except in the special fiber, which is an endless chain of rational curves. There is only one non-trivial 2-block partition of the relevant modular graph, so we need only specify integers N_u and N_l. In this situation, the inequalities above pick out the subscheme S_{N_u,N_l} illustrated below; note that this scheme includes all Gieseker bundles on smooth curves, but only those strata in the special fiber which "lie between" the fixed points singled out by N_u and N_l



Remark 3.12. Roughly speaking, the spaces S_{N_u,N_l} are obtained by deleting the "infinite tails" of A_d, which classify bundles where the splitting degree becomes large. We shall make this more precise below.

3.5. **Finite-Type Substacks and Fixed Point Strata.** It is intuitively clear that there must be some relationship between the finite-type spaces S_{N_u,N_l} defined above and the fixed points of the groups $\mathcal{G}_R$: The inequalitities in Definition 3.10, in essence, forbid us from using Gieseker bubbles to shift degree across the nodes. In this subsection, we make these ideas precise.

3.5.1. *Deformations which Smooth Gieseker Bubbles.* Fix a non-trivial 2-block partition R of V_{γ_o} and consider $F_R^{\underline{n}}$, the $\underline{n}$-th connected component of the fixed point locus F_R of $\mathcal{G}_R$. The bundles in the locus $F_R^{\underline{n}}$ have Gieseker bubbles at the nodes labelled by the splitting edges in E_R^{split}.

A bundle classified by $F_R^{\underline{n}}$ has Gieseker bubbles at each node labelled by one of the splitting edges in E_R^{split}; each such Gieseker bubble has two nodes, both of which stabilize to the same node. Let $D_R^{\underline{n}}$ be the subspace of A_d which classifies bundles obtained by smooth at most one of the nodes on each of the Gieseker bubbles labelled by E_R^{split}.

The condition we've used to define $D_R^{\underline{n}}$ can be stated in terms of labelled deformations of modular graphs, so $D_R^{\underline{n}}$ is a union of strata labelled by such.

Example 3.13. In the special case of Example 1.12, where there is only one non-trivial two-block partition, the locus D^n containing the fixed point set $F^n = \mathrm{pt}_n$ is

$$D^n = (U_n)_0 = \operatorname{Spec} \frac{\mathbb{C}[z_n, w_n]}{\langle z_n w_n \rangle} =$$  pt_n.

This is a pair of affine lines meeting at a common origin pt_n.

The $(\mathbb{C}^\times)^V$ action on A_d respects the modular graph stratification, so the action of $\mathcal{G}_R$ on A_d restricts to an action of $\mathcal{G}_R$ on $D^{\underline{n}}_R$. Since $\mathcal{G}_R$ acts nontrivially on all the strata in $D^{\underline{n}}_R \setminus F^{\underline{n}}_R$, we may think of these strata as flowing "towards" or "away" from the fixed point locus $F^{\underline{n}}_R$.

Definition 3.14. $Z^{\underline{n}}_R$ is the algebraic subspace of $D^{\underline{n}}_R$ for which the weights of $\mathcal{G}_R$ on the conormal bundle $\overline{N}_{F^{\underline{n}}_R/Z^{\underline{n}}_R}$ are all *non-negative.*

Likewise, $W^{\underline{n}}_R \subset D^{\underline{n}}_R$ is the algebraic subspace for which the weights of $\mathcal{G}_R$ on the conormal bundle $\overline{N}_{F^{\underline{n}}_R/W^{\underline{n}}_R}$ are all *non-positive.*

Example 3.15. In the special case of Example 1.12,

$$Z^n = \operatorname{Spec} \mathbb{C}[z_n] =$$  pt_n

and

$$W^n = \operatorname{Spec} \mathbb{C}[w_n] =$$  pt_n.

In the example, above Z^n and W^n are the total spaces of vector bundles (of rank 1) over the fixed point locus F^n. An analogous statement is generally true.

Definition 3.16. The *projection map* $\eta_Z : Z^{\underline{n}}_R \to F^{\underline{n}}_R$ sends a Gieseker bundle $(C, \mathcal{P})$ to the Gieseker bundle obtained by

1. creating a Gieseker bubble at every node σ_e labelled by an $e \in E^{split}_R$ for which the modification $m : C \to \Sigma$ is trivial, and
2. for each such node σ_e, twisting the bundle on the curve component on one side of σ_e by the divisor $-\sigma_e$. Which side one twists at is determined by the requirement that the resulting multidegree land in the $\underline{n}$-th fixed point locus.

Similarly, we have a projection map $\eta_W : W^{\underline{n}}_R \to F^{\underline{n}}_R$.

Proposition 3.17. *$Z^{\underline{n}}_R$ is the total space of the conormal bundle $\overline{N}_{F^{\underline{n}}_R/Z^{\underline{n}}_R}$. Likewise, $W^{\underline{n}}_R$ is the total space of the conormal bundle $\overline{N}_{F^{\underline{n}}_R/W^{\underline{n}}_R}$.*

Proof. The fiber of the projection map η_Z over a given Gieseker bundle in $F^{\underline{n}}_R$ is the stack of bundles on the union of all Gieseker $\mathbb{P}^1$'s and points which can arise as the preimages under the modification map of nodes in Σ_b labelled by edges $e \in E^{split}_R$. The stack of such bundles on a point is a copy of $\mathbb{C}^\times$, and the stack of such bundles on a Gieseker $\mathbb{P}^1$ is a pt. Thus, every fiber of η_z is a copy of $\mathbb{A}^{|E_R^{split}|}$; the Gieseker strata give a toric decomposition.

The proof for η_w is identical. ☐

Corollary 3.18. *$Z_R^{\underline{n}}$ and $W_R^{\underline{n}}$ are smooth.*

Proof. $Z_R^{\underline{n}}$ is the total space of a vector bundle on the fixed point locus $F_R^{\underline{n}}$. ☐

This fact will be important in our proof of the main theorem.

3.5.2. *An Alternate Description of S_{N_u,N_l}.* Pick a partition $R_0 \in NT2B(\gamma_o)$ and define $M_u(R)$ and $M_l(R)$ by

$$M_u(R) = \begin{cases} N_u(R)+1 & \text{if } R = R_0 \\ N_u(R) & \text{otherwise} \end{cases} \qquad M_l(R) = \begin{cases} N_l(R)-1 & \text{if } R = R_0 \\ N_l(R) & \text{otherwise.} \end{cases}$$

Proposition 3.19. *S_{N_u,N_l} is the complement of $W_R^{N_l(R)}$ in S_{N_u,M_l}:*

$$S_{N_u,N_l} = S_{N_u,M_l} \setminus W_R^{N_l(R)}.$$

Likewise, S_{N_u,N_l} is the complement of $Z_R^{N_u(R)}$ in S_{M_u,N_l}:

$$S_{N_u,N_l} = S_{M_u,N_l} \setminus Z_R^{N_u(R)}.$$

Proof. This is an elementary observation: Deformations which land in $T_R^Z(N_u)$ shift all of the degree 1's from the Gieseker bubbles to the components in the block V^+ of R. The inequality

$$d_+(\gamma, R) \geq d + N_l(R) - k(R) + 1$$

says precisely that these deformations are forbidden. ☐

The proposition implies that S_{N_u,N_l} is obtained by deleting the "infinite tails" of A_d, generalizing the picture given in the examples involving Gieseker bundles on curves of genus 0 with 4 marked points. More precisely:

Definition 3.20. Let $N_u = (N_u(R))$ and $N_l = (N_l(R))$ be as in Definition 3.10. For any partition $R \in NT2B(\gamma_o)$, consider the following algebraic subspaces of A_d:

$$T_R^Z(N_u) = \bigcup_{n > N_u(R)} Z_R^n \qquad \text{and} \qquad T_R^W(N_l) = \bigcup_{n < N_l(R)} W_R^n.$$

These loci are the *infinite tails* associated to the partition R and the indices N_u and N_l.

Proposition 3.21. *The space S_{N_u,N_l} is obtained as the complement*

$$S_{N_u,N_l} = A \setminus \bigcup_{R \in NP_2(\gamma_o)} T_R^Z(N_u) \sqcup T_R^W(N_l).$$

of the union of the infinite tails.

3.6. **Proper Substacks.** Obviously, the subspaces S_{N_u,N_l} get smaller as we make N_u and N_l closer together. We now consider the extremal case where

$$N_u(R) = N_l(R) + 1 = N(R)$$

for all $R \in NT2B(\gamma_o)$. The corresponding algebraic subspace is denoted

$$S_N = S_{(N(R)),(N(R)-1)}.$$

Lemma 3.22. *The stabilizer group of any Gieseker bundle classified by S_N is $\mathbb{C}^\times_\Delta$.*

Proof. The inequalities in the definition eliminate from S_N any stratum $A_{\gamma,\underline{d}}$ which, for any non-trivial 2-block partion R, has a Gieseker bubble on every node connecting components labelled vertices in different blocks of R. This means that any two components are connected by a path which does not pass through any unstable $\mathbb{P}^1$'s. It follows that the stabilizer group of any point in $S_{\underline{n}}$ is the diagonal $\mathbb{C}^\times_\Delta$. □

Suppose that choose a particular $v_0 \in V$ and thereby split $(\mathbb{C}^\times)^V \simeq \mathbb{C}^\times_\Delta \times (\mathbb{C}^\times)^{V\setminus\{v_0\}}$. The lemma above says that $(\mathbb{C}^\times)^{V\setminus\{v_0\}}$ acts freely on S_N, allowing us to forget all but one of the trivializations, by fixing them in reference to t_{v_0}. More precisely, the quotient stack $[S_N/(\mathbb{C}^\times)^V]$ can be written as

$$[S_N/(\mathbb{C}^\times)^V] \simeq Q_N \times [\mathrm{pt}/\mathbb{C}^\times_\Delta],$$

where Q_N is the stack which classifies degree d Gieseker bundles equipped with a single trivialization t_{v_0} (or equivalently, the coarse moduli space f degree d Gieseker bundles) obeying the inequalities from the definition.

Proposition 3.23. *The coarse moduli space Q_N defined above is proper, relative to B.*

To prove this, we shall need some facts about the multidegrees of bundles which can occur in the special fiber of a family on the Spec of a discrete valuation ring. Let R be a discrete valuation ring, with fraction field K, and associated disc $D = \mathrm{Spec}(R)$ and punctured disc $D^\times = \mathrm{Spec}(K)$.

Definition 3.24. We will say that two principal $\mathbb{C}^\times$-bundles $\mathcal{P}_o$ and $\mathcal{P}'_o$ on the special fiber Σ_o are *fiber twists* of one another if $\mathcal{P}_o \simeq \mathcal{P}'_o \times_{\mathbb{C}^\times} \mathcal{T}$, where $\mathcal{T}$ is the restriction to Σ_o of the principal bundle associated to the divisor $\sum_{v\in V_{\gamma_o}} n_v \Sigma_{o,v}$ on Σ.

Suppose we are given a family of curves Σ/D on the disc, with smooth generic fiber, and a family of $\mathbb{C}^\times$-bundles $\mathcal{P}/(\Sigma|_{D^\times})$ over the punctured disc $D^\times$. If the family $\mathcal{P}$ extends to D, with special fiber $\mathcal{P}_o$, then it can also be extended to D with special fiber $\mathcal{P}'_o$ any twist of $\mathcal{P}_o$. When the generic fiber is nodal, one can repeat this story by focusing attention on each smooth component.

The set of multi-degrees of twists of the trivial bundle may be identified as follows: Consider the intersection matrix $k = (k_{vv'})$ of Σ_o. If $v \neq v'$, then $k_{vv'} = |\Sigma_{o,v} \cap \Sigma_{o,v'}|$, the number of nodes common to both curves. If $v = v'$, $k_{vv} = -|\Sigma_{o,v} \cap \overline{\Sigma_o \setminus \Sigma_{o,v}}|$, the number of nodes where $\Sigma_{o,v}$ meets the closure of its complement.

Lemma 3.25 ([Cap08])**.** *The set of multidegrees of twists of the trivial bundle is the lattice $\Lambda_{\Sigma_o} \subset \mathbb{Z}^{V_{\gamma_o}}$ generated by the columns of the intersection matrix k.*

Proof. This follows from the fact that the degree $\deg_{\Sigma_{o,v}} \mathcal{O}_\Sigma(\Sigma_{o,v'})$ is equal to $k_{vv'}$. □

Proof of Prop. 3.23. Q_N is clearly of finite type.

We show that the valuative criteria for completeness and separability are satisfied. Let R be a discrete valuation ring, with fraction field K, and associated disc $D = \operatorname{Spec}(R)$ and punctured disc $D^\times = \operatorname{Spec}(K)$. We claim that, given any family $(C, \mathcal{P})$ of Gieseker bundles on $D^\times$ which satisfy the inequalities from the definition, there exists a unique extension of $(C, \mathcal{P})$ to D whose special fiber satisfies the inequalities.

The stack $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is complete, so there is some extension of $\mathcal{P}$ to the entire disc D. The set of all possible extensions consists of all twists of a certain bundle $\mathcal{P}_o$ on the special fiber of C, and clearly at least one of these twists will satisfy the inequalities from the definition. Moreover, only one of these twists can lie in Q_N, so Q_N is separated. This follows directly from the characterization of multidegrees of twists of the trivial bundle in Lemma 3.25: A non-trivial twist will always shift the degree on some component of the special fiber by a multiple of $k(R)$. In the special case $N_u(R) = N_l(R) + 1$, this leads to a violation of the inequalities. Thus, any non-trivial twist of a Gieseker bundle in Q_N lies outside of $Q_{\underline{n}}$. □

4. Admissible Classes

In this section, we introduce the admissible classes, and estimate the weights of their fibers over fixed points $F_R^{\underline{n}}$ as functions of $\underline{n}$.

4.1. **Definitions.** The definitions given here are relative versions of the ones given in [TW03].

Recall that $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ carries a universal curve C with universal marked points $\sigma_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to C$, and a universal bundle $p : \mathcal{P} \to C$. The universal bundle may be thought of equivalently as a morphism $\phi : C \to [\mathrm{pt}/\mathbb{C}^\times]$, and the universal marked points give rise to evaluation maps $\mathrm{ev}_i : \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \to [\mathrm{pt}/\mathbb{C}^\times]$.

For any finite-dimensional representation V of $\mathbb{C}^\times$, let $\mathcal{V} = \phi^* V$ be the vector bundle on $\Sigma_{g,I}$ associated to V by the universal bundle.

Definition 4.1. The *evaluation class* $\mathrm{ev}_i^*[V]$ is the K-theory class on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ represented by the vector bundles $\sigma_i^*\mathcal{V}$.

A *descendant class* is a K-theory class of the form $\mathrm{ev}_i^*[V] \otimes [T_i^{\otimes n_i}]$, where $T_i = \sigma_i^* T_\pi^*$ is the pullback of the relative cotangent bundle of the morphism $\pi : C \to \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ and n_i is a positive integer.

Remark 4.2. These are K-theoretic gravitational descendents. Their cohomological analogues, the ψ-classes, are the images of these classes under the Chern character: $\mathrm{ch}([T_i]) = \exp(\psi_i)$.

Definition 4.3. The *Dolbeault index class I_V of a representation V* is the topological K-theory classes associated to the complexes $R\pi_*\mathcal{V}$ of sheaves on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$.

The *determinant of cohomology of V* is the line bundle $\det I_V = \det R\pi_*\mathcal{V}$.

Definition 4.4. A line bundle $\mathcal{L}$ on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is *admissible* if it is topologically isomorphic to a positive (possibly fractional) power of the dual of the determinant of cohomology of the standard representation $\mathbb{C}$ of $\mathbb{C}^\times$:

$$\mathcal{L}^{-1} \simeq (\det R\pi_*\phi^*\mathbb{C})^{\otimes q}$$

for some positive rational number q.

Definition 4.5. An *admissible complex* α on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$ is the tensor product of an admissible line bundle $\mathcal{L}$ with any number of Dolbeault index and evaluation/descendant classes.

$$\alpha = \mathcal{L} \bigotimes \otimes_a (R\pi_*\mathcal{V}_a) \bigotimes \otimes_i (\mathrm{ev}_i^* V_i \otimes T_i^{\otimes n_i}).$$

An *admissible class* is a topological K-theory class represented by sums of admissible complexes.

4.2. **Weight Estimates in the Local Atlas.** The local atlas A_d has universal curves and bundles as well, and one can define the admissible classes associated to these universal structures. These admissible classes, so defined, agree with the pullback of admissible classes from the stack $\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$. We will use the same notation for classes on A_d that we used for classes on $\widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times])$.

Admissible complexes are complexes of coherent sheaves on A_d, so we can represent them as complexes $\mathcal{V}^\bullet$ of vector bundles. If $f \in F_R^{\underline{n}}$ is a $\mathcal{G}_R$-fixed point, then the fiber $\mathcal{V}_f^i$ of $\mathcal{V}^i$ at f is a $\mathcal{G}_R$-representation. The weights of these representations, being both discrete-valued and continuous in f, depend only on the connected component $F_R^{\underline{n}}$. Our goal in this section is to estimate the $\mathcal{G}_R$-weights of the fixed-point fibers of $\mathcal{V}^i$ as functions of $\underline{n}$.

Proposition 4.6. *For any 2-block partition $R \in NT2B(\gamma_o)$,*

1. *The $\mathcal{G}_R$-weights of the fixed point fibers of the evaluation and descendant classes are bounded functions of $\underline{n}$.*

(2) *The index complex* $R\pi_*\mathcal{V}$ *is quasi-equivalent to a complex* $L^\bullet$ *of vector bundles, and the* $\mathcal{G}_R$*-weights of the fixed point fibers of each bundle* L^i *in this complex are bounded functions of* $\underline{n}$*.*

(3) *The* $\mathcal{G}_R$*-weights of fixed-point fibers of admissible line bundles* $\mathcal{L}$ *grow linearly with* $\underline{n}$*, with positive coefficient.*

The remainder of this section is devoted to proving this proposition.

Let $f \in F_R^{\underline{n}}$ be a fixed point of $\mathcal{G}_R \subset \mathcal{G}$, represented by $(C_f, \sigma_{f,i}, \mathcal{P}_f, \{t_{v,f}\})$. We denote the modular graph of $(C_f, \sigma_{f,i})$ by γ.

Lemma 4.7. *Let* $\mathbb{C}_\lambda$ *be the irreducible* $\mathbb{C}^\times$*-representation of weight* λ*, and let* $\mathcal{V}_\lambda|_{C_f} = \mathcal{P}_f \times_{\mathbb{C}^\times} \mathbb{C}_\lambda$ *be the associated bundle on* C_f*. Recall that* $\mathcal{G}_R$ *is a product* $\Pi_r \mathbb{C}^\times_{\Delta^r}$*. Let* U *be an open subset of* C_f *which is entirely contained in one of the components* $C_{v,f}$ *of* C_f*. If* $C_{v,f}$ *is one of the components of the block* V^r_γ*, then* $\mathbb{C}^\times_{\Delta^r}$ *acts on* $\Gamma(U, \mathcal{V}_\lambda|_{C_f})$ *with weight* $-\lambda$*. Otherwise,* $\mathbb{C}^\times_{\Delta^r}$ *acts on* $\Gamma(U, \mathcal{V}_\lambda|_{C_f})$ *with weight* 0*.*

Proof. f is a $\mathcal{G}_R$-fixed point, so if we scale the trivializations t_v $(v \in V^r)$ by $g_r \in \mathbb{C}^\times_{\Delta^r}$, we can compensate by rescaling the bundle's fibers on the appropriate components of C_f by g_r^{-1}. This rescales the local sections of the associated bundles $\mathcal{V}_\lambda|_{C_f}$ by the appropriate power $g_r^{-\lambda}$. □

4.2.1. *Evaluation and Descendant Classes.* The fixed-point fiber of an evaluation bundle $\mathrm{ev}_i^* V = \sigma_i^*\mathcal{V}$ is simply the fiber of $\mathcal{V}$ at the image $\sigma_i(f)$ of the fixed point inside C_f. Any $\mathbb{C}^\times$-representation V is a direct sum $\oplus_a \mathbb{C}_{\lambda_a}$ of irreps, so Lemma 4.7 implies that the $\mathcal{G}_R$-weights of the fixed-point fibers of the evaluation bundle are constant functions of $\underline{n}$.

Similarly, $\mathcal{G}_R$ acts trivially on the stable components of C_f (although it rotates the Gieseker bubbles), so the same reasoning show that weights of the fixed-point fibers of the descendant class $\sigma_i^*(\mathcal{V} \otimes (T_\pi^*)^{\otimes k})$ are constant in $\underline{n}$.

4.2.2. *Index Classes.* The morphism $\pi : C \to A_d$ is projective, the atlas A_d is locally Noetherian, and the associated vector bundle $\mathcal{V}$ on C (being a locally-free sheaf) is flat over A_d. It follows (Hartshorne III.12 [Har77], p. 282-4) that, locally on A_d, $R\pi_*\mathcal{V}$ is quasi-equivalent to a complex $L^\bullet$ of finitely-generated locally-free sheaves. These locally-free sheaves are generated by local sections $\{s_\alpha\}$ of a Cech resolution of $\mathcal{V}$; each such section s_α represents a chosen generator of the sheaves $R^i f_*\mathcal{V}$.

The fixed-point fibers of the L^i are spanned by the images of the generators s_α in the fibers. Such images are represented by local sections of $\mathcal{V}|_{C_f}$, so Lemma 4.7 above implies that the $\mathcal{G}_R$-weights on the fixed point fibers don't depend on $\underline{n}$.

4.2.3. *Admissible Line Bundles.* Any admissible line bundle $\mathcal{L}$ is topologically a positive (possibly fractional) power $(\mathcal{L})^q$ of the determinant class $\mathcal{L}_1 = \det^{-1} R\pi_* \mathcal{V}_1$ associated to the standard representation $\mathbb{C}_1$ of $\mathbb{C}^\times$. The $\mathcal{G}_R$-fixed point weights of $(\mathcal{L}_1)^{\otimes q}$ are linear in q, so it's enough to compute them for $\mathcal{L}_1$.

Dual and determinant are functorial operations, so, as elements of $K_{\mathcal{G}_R}(f)$,

$$[\mathcal{L}_1|_f] = (\det[R\Gamma(C_f, \mathcal{V}_1|_{C_f})])^{-1}.$$

These K-theory classes are topological invariants, so they are unchanged by deformations within $F_R^{\underline{n}}$, so we may assume C_f has only those nodes that it must have represents a $\mathcal{G}_R$-fixed point (i.e., those nodes which create the required Gieseker bubbles). Furthermore, these characters are necessarily polynomial in the components n_v of $\underline{n}$, so it's enough to compute them when $n_v \gg 0$ for all $v \in V_\gamma$. In this situation, there are no higher cohomologies, and the global sections in $\Gamma(C_v, \mathcal{V}_1|_{C_f})$ are specified by giving sections of $\mathcal{V}_1|_{C_f}$ on the irreducible components $C_{v,f}$ of C_f. (The sections on the Gieseker bubbles are determined by continuity at the nodes, since the space of possible such sections is two-dimensional.) The weights of $\mathcal{G}_R$ on these sections were computed in Lemma 4.7.

$$[R\Gamma(C_f, \mathcal{V}_1|_{C_f})] = \sum_{r \in V_\gamma} (n_r + 1 - g_r) t_r^{-1}.$$

Thus, we have

Proposition 4.8.

$$[\mathcal{L}|_f] = \Pi_{r \in V_\gamma} t_r^{q(n_r+1-g_r)}.$$

As promised, the fixed point weights grow linearly in $\underline{n} = (n_r)$, with positive coefficient.

5. Gromov-Witten Invariants for $[\mathrm{pt}/\mathbb{C}^\times]$

Definition 5.1. Let $[V_i] \in K([\mathrm{pt}/\mathbb{C}^\times])$ be a collection of K-theory classes, indexed by $i \in I$. Let $\mathcal{L}$ be an admissible line bundle, and let $\otimes_a I_{V_a}^{\otimes n_a}$ be a collection of index classes. Set

$$h = \mathcal{L} \otimes \otimes_a I_{V_a}^{\otimes n_a}.$$

The *h-twisted Gromov-Witten invariant* $\langle V_1, ..., V_{|I|}\rangle_h$ *associated to the collection* $\{V_i\}$ is the Euler characteristic of the admissible complex

$$h \bigotimes \otimes_i \mathrm{ev}_i^* V_i = \mathcal{L} \bigotimes \otimes_a (R\pi_* \mathcal{V}_{\lambda_a}) \bigotimes \otimes_i \mathrm{ev}_i^* V_i$$

The h-twisted invariants for descendant classes are defined similarly.

5.1. **Statement of The Main Theorem.** The Euler characteristic of a complex α on $\widetilde{\mathcal{M}}_{g,I}(\mathrm{pt}/\mathbb{C}^\times)$ is the Euler characteristic of the right-derived pushforward $R(F_{\mathcal{P}})_*\alpha$. The fibers of F are not proper, so it is not obvious that these invariants are well-defined.

The remainder of this section is devoted to proving that the following theorem, which implies that the K-theory class

$$F_{\mathcal{P}*}[\alpha] := [RF_{\mathcal{P}*}\alpha] = \sum_i (-1)^i [I^i(RF_*\alpha)]$$

is a finite sum, hence well-defined. (Here I is any locally free resolution of $RF_*\alpha$. Such resolutions exist because $\overline{\mathcal{M}}_{g,I}$ is smooth and projective. The K-theory class is independent of which resolution we choose.)

Theorem 5.2. *The derived pushforward $R^\bullet F_*\alpha$ of any admissible complex $\alpha^\bullet$ is coherent.*

5.2. **Proof of the Main Theorem.** Coherence is a local property, so we can check the coherence of $RF_*\alpha$ in our favorite atlas $A = \sqcup_{d\in\mathbb{Z}} A_d$.

$$\begin{tikzcd}
A \arrow[r, "q"] \arrow[dr, "f"'] & \widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)} \arrow[r] \arrow[d, "F"] & \widetilde{\mathcal{M}}_{g,I}([\mathrm{pt}/\mathbb{C}^\times]) \arrow[d, "F"] \\
 & B \arrow[r] & \overline{\mathcal{M}}_{g,I}
\end{tikzcd}$$

B is affine and $q : A \to \widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$ presents $\widetilde{\mathcal{M}}_{(\Sigma,\sigma_i)}$ as a quotient by $(\mathbb{C}^\times)^V$, so proving coherence amounts to proving that the $(\mathbb{C}^\times)^V$ invariants in the derived global sections $R\Gamma(A,\alpha)$ are finitely-generated.

We'll prove this in two steps: First, we'll work at fixed total degree d, showing that the $(\mathbb{C}^\times)^V$ invariants in the derived global sections $R\Gamma(A_d,\alpha)$ are coherent. Then, we'll show that these invariants vanish for all but finitely many d, so that the sum

$$[R\Gamma(A,\alpha)]^{(\mathbb{C}^\times)^V} = \sum_{d\in\mathbb{Z}} [R\Gamma(A_d,\alpha)]^{(\mathbb{C}^\times)^V}$$

is well-defined.

5.2.1. *Coherence on A_d.* There are two cases: $|V_{\gamma_o}| = 1$ and $|V_{\gamma_o}| \geq 2$.

The first case is essentially trivial. If $|V_{\gamma_o}| = 1$, then, as we have observed, A_d is the compactified Jacobian, proper over B. In this case $R\Gamma(A_d,\alpha)$ is finitely-generated, as is its submodule of $(\mathbb{C}^\times)^V \simeq \mathbb{C}^\times$-invariants.

The second case – $|V_{\gamma_o}| \geq 2$ – is more complicated, because A_d is not proper over B. It has infinitely many finite-type strata. We would like to show that most of these strata do not contribute to $(\mathbb{C}^\times)^V$-invariants to $R\Gamma(A_d,\alpha)$.

Recall that we can obtain A_d as the limit

$$A_d = \lim_{N_u(R)\to\infty, N_l(R)\to -infty} S_{N_u,N_l}.$$

The following statement about local cohomology allows us to reduce the question of finite-generation on A_d to the question of finite-generation on any S_{N_u,N_l}.

Lemma 5.3. *Let $\mathcal{V}$ be a finite rank $(\mathbb{C}^\times)^V$-equivariant vector bundle on A_d for which, for all non-trivial two-block partitions R, the $\mathcal{G}_R$-weights of the fibers of $\mathcal{V}$ over a $\mathcal{G}_R$-fixed point $F_R^{\underline{n}}$ are linear functions of $\underline{n}$, with positive coefficients on the linear terms.*

Fix a partition R and multidegree bounds N_u and N_l, and abbreviate

$$S = S_{N_u,N_l} \qquad Z = Z^{N_u(R)-1}(R)_R \qquad W = W_R^{N_l(R)+1}.$$

The $(\mathbb{C}^\times)^V$-invariants in the local cohomology groups

$$R^p\Gamma_Z(S,\mathcal{V}) \qquad \textit{and} \qquad R^p\Gamma_W(S,\mathcal{V})$$

are finitely generated.

Moreover, these local cohomologies vanish if $N_u(R) \gg 0$ and $N_l(R) \ll 0$ for all R.

Proof. The argument for W has the same form as the argument for Z, so we focus on the latter case.

The sheaf $\mathcal{V}$ is torsion-free, so the local cohomology sheaf $\Gamma_Z(\mathcal{V})$ vanishes. In fact, because Z is a closed connected subvariety of S (of codimension q), the only non-zero local cohomology sheaf is $R^q\Gamma_Z(\mathcal{V})$. The local-to-global spectral sequence, together with the exactness of the $(\mathbb{C}^\times)^V$-invariants functor, implies that the vanishing of the $\mathcal{G}_R$-invariants in the local cohomology groups $R^i\Gamma_Z(S,\mathcal{V})$ follows from the vanishing of the $\mathbb{C}_R^\times$-invariants in $R^p\Gamma(A, R^q\Gamma_Z(\mathcal{V}))$. The vanishing of the latter invariants follows (via the filtration spectral sequence) from the vanishing of the $\mathcal{G}_R$-invariants in the cohomology groups $R^j\Gamma(Z, \mathcal{V}\otimes \operatorname{Sym} N_{Z/S})$.

Z is the total space of a vector bundle over the fixed point locus $F = F_R^{N_u(R)-1}$, so (taking global sections over the fibers, we see that

$$R^i\Gamma(Z,\mathcal{V}\otimes \operatorname{Sym} N_{Z/S}) = R^i\Gamma(F, \mathcal{V}\otimes \operatorname{Sym} N_{Z/S} \otimes \operatorname{Sym} \overline{N}_{F/Z}).$$

The weight spaces in the two Sym's in the RHS above are finitely generated, and vanish for negative weights. Since $\mathcal{V}$ is finite rank, it follows that the $(\mathbb{C}^\times)^V$-invariants in the RHS are finitely-generated. Moreover, because we have assumed that the fixed-point weights of $\mathcal{V}$ grow linearly, with positive coefficients, the $(\mathbb{C}^\times)^V$-invariants in the RHS vanish if N_u and N_l lie outside some finite range. □

We saw in Section 4 that an admissible complex can be represented as complex of bundles $\mathcal{V}$ which satisfy the conditions of the lemma. (The evaluation, descendant, and index classes are all bounded, and the admissible line $\mathcal{L}$ supplies the linear growth.)

Corollary 5.4. *There exist* N_u *and* N_l *such that*

$$R\Gamma(A_d,\alpha)^{(\mathbb{C}^\times)^V} = R\Gamma(S_{N_u,N_l},\alpha)^{(\mathbb{C}^\times)^V}.$$

Proof. The local cohomology groups in the lemma measure how $R\Gamma(S_{N_u,N_l},\alpha)$ changes if we increase $N_u(R)$ or decrease $N_l(R)$ by 1. Since the invariants in these local cohomologies vanish outside some finite range of N_u and N_l the limit

$$\lim_{N_u(R)\to\infty, N_l(R)\to-\infty} R\Gamma(S_{N_u,N_l},\alpha)^{(\mathbb{C}^\times)^V}$$

stabilizes. □

The finite-generation of invariants in the local cohomology also lets us delete strata from S_{N_u,N_l}. This may alter the $(\mathbb{C}^\times)^V$-invariants, but because the local-cohomologies are finitely-generated, it will not alter the fact of their finite-generation. So, we can check finite-generation by reducing to the "minimal" subspace S_N $(= S_{N,N-1})$. But, by Proposition 3.23, the invariants in $R\Gamma(S_N,\alpha)$ are the invariants in $R\Gamma(Q_n,\alpha)$. Q_N is coherent, so the latter invariants are automatically finitely generated.

Corollary 5.5. *The* $(\mathbb{C}^\times)^V$*-invariants in* $R\Gamma(A_d,\alpha)$ *are finitely-generated.*

5.2.2. *The Sum over d.* Now we would like to show that the $(\mathbb{C}^\times)^V$-invariants in $R\Gamma(A_d,\alpha)$ vanish, for all but finitely many d.

The diagonal subgroup $\mathbb{C}^\times_\Delta \subset (\mathbb{C}^\times)^V$ fixes all of A_d, so the global sections $R\Gamma(A_d,\alpha)$ form a complex of $\mathbb{C}^\times_\Delta$-representations. The $\mathbb{C}^\times_\Delta$-weight spaces in this complex are non-zero for only finitely many weights. The width of this range depends on the class α; however, the growth of the upper and lower bounds of the range depend only on the admissible line bundle factor $\mathcal{L}$ in α. In particular, the upper and lower bounds grow linearly in d, with positive coefficient. (This is a consequence of Proposition 4.8; the weights of the diagonal $\mathbb{C}^\times_\Delta$ are proportional to $q\sum_r(n_r+1-g_r) = d+\text{constant}$.) Vanishing for all but finitely many d follows.

6. Towards Gromov-Witten Invariants for $[X/\mathbb{C}^\times]$

6.1. **Definitions.** Recall that $[X/\mathbb{C}^\times]$ is, by definition, the fibered category whose objects are triplets $(B,\mathcal{P},s)$ consisting of a test scheme B, a principal $\mathbb{C}^\times$-bundle $p:\mathcal{P}\to B$, and a $\mathbb{C}^\times$-equivariant morphism $s:\mathcal{P}\to X$. Morphisms between such pairs are Cartesian diagrams

$$\begin{tikzcd}
\mathcal{P} \arrow[r, "f"] \arrow[d, "p"] & \mathcal{P}' \arrow[d, "p'"] \\
B \arrow[r] & B'
\end{tikzcd}$$

such that $s = s'\circ f$.

The upshot of this definition is that the natural map $\rho : X \to [X/\mathbb{C}^\times]$ is a principal $\mathbb{C}^\times$-bundle, and any map $\phi : C \to [X/\mathbb{C}^\times]$ gives rise to a pullback diagram

$$\begin{tikzcd}
\mathcal{P} \arrow[r, "s"] \arrow[d, "p"] & X \arrow[d, "\rho"] \\
C \arrow[r, "\phi"] & {[X/\mathbb{C}^\times]}
\end{tikzcd}$$

Thus maps to $[X/\mathbb{C}^\times]$ *are* principal $\mathbb{C}^\times$-bundles, together with a section s of the associated fiber bundle $\mathcal{P} \times_{\mathbb{C}^\times} X$.

There is a natural notion of degree for such maps. Define the homology of $[X/\mathbb{C}^\times]$ by the equation $H_n([X/\mathbb{C}^\times]) = H^{\mathbb{C}^\times}_{n+\dim(\mathbb{C}^\times)}(X)$, so that the image of the $\mathbb{C}^\times$-equivariant fundamental class $[\mathcal{P}]^\times_{\mathbb{C}}$ is the usual fundamental class $[C]$. We will say that a map $\phi : C \to [X/\mathbb{C}^\times]$ has *degree* $\beta \in H^{\mathbb{C}^\times}_{2+\dim(\mathbb{C}^\times)}(X) = H_2([X/\mathbb{C}^\times])$ if the pushforward $s_*[\mathcal{P}]_{\mathbb{C}^\times} = \beta$. For a map to $\mathrm{pt}/\mathbb{C}^\times$, this notion of degree is equivalent to the usual definition of a bundle's degree via the first Chern class.

Definition 6.1. A *Gieseker map from C to $[X/\mathbb{C}^\times]$* is a triplet $((C, \sigma_i), \mathcal{P}, s)$ consisting of:

1. a prestable marked curve (C, σ_i),
2. a principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$, and
3. A section $s : \mathcal{P} \to X$,

such that

1. $\mathcal{P}$ has degree 0 on any irreducible rational component of C which has one node and one marked point.
2. $\mathcal{P}$ has either degree 0 or degree 1 on any unstable rational component of C which has two nodes.
3. s is non-trivial on any unstable component on which $\mathcal{P}$ has degree 0.

A Gieseker map has degree $\beta \in H_2([X/\mathbb{C}^\times])$ if $s_*[\mathfrak{c}^*\mathcal{P}]_{\mathbb{C}^\times} = \beta$.

We denote by $\widetilde{\mathcal{M}}_{g,I}([X/\mathbb{C}^\times])$ the fibered category of Gieseker maps to $[X/\mathbb{C}^\times]$ from stable marked curves of type (g, I). Its connected components are labelled by the degree $\beta \in H_2([X/\mathbb{C}^\times])$; we denote them by $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$.

It is clear from the definition that there is a forgetful map

$$F_s : \widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times]) \to \widetilde{\mathcal{M}}_{g,I,ft_*\beta}(\mathrm{pt}/\mathbb{C}^\times),$$

where $ft_*\beta$ is the degree obtained from the homomorphism $ft_* : H_2([X/\mathbb{C}^\times]) \to H_2([\mathrm{pt}/\mathbb{C}^\times])$. We forget the section s and we obtain a Gieseker map to $[\mathrm{pt}/\mathbb{C}^\times]$ by contracting any rational components which carry trivial bundles.

Remark 6.2. This definition is inspired by Kontsevich's definition of stable maps. Sections $s : \mathcal{P} \to X$ are locally maps from C to X. Thus, sections can degenerate in families in exactly the same way that maps to X do, by developing singularities. We cure these singularities by bubbling where the singularity occurs.

Theorem 6.3. *F_s is proper and Deligne-Mumford.*

Proof. In essence, the result follows from the fact that $[X/\mathbb{C}^\times] \to [\mathrm{pt}/\mathbb{C}^\times]$ is proper and representable. We make the argument precise by proving valuative criteria for completeness & separability. Let R be a discrete valuation ring (over $\mathbb{C}$), with fraction field K and consider the associated disc $D = \mathrm{Spec}(R)$ and punctured disc $D^\times = \mathrm{Spec}(K)$.

Let us suppose that we have a family $(C, \sigma_i, \mathcal{P}, s)$ over $D^\times$, and an extension $(Z, z_i, \mathcal{R})$ of $F_s(C, \sigma_i, \mathcal{P})$ to D. (Any required base changes will be subsumed in the notation.)

Completeness:

We want to exhibit an extension $(Y, y_i, \mathcal{Q}, t)$ of $(C, \sigma_i, \mathcal{P}, s)$ to D such that $F_s(Y, y_i, \mathcal{Q}, t) = (Z, z_i, \mathcal{R})$.

First, we extend the family (C, σ_i) to D. This may require base change, and is an easy consequence of the existence of nodal reduction [HM98]. We denote the extension by (Y', y_i'). Y' comes equipped with a contraction map $c : Y' \to Z$. The marked We denote by $\mathcal{Q}'$ the pullback $c^*\mathcal{R}$; note that $\mathcal{Q}'$ is trivial on components collapsted by c.

Given Y', the graph of s gives us an embedding $j : C \to X_{\mathcal{Q}'}$, where $X_{\mathcal{Q}'}$ is the associated bundle $\mathcal{Q}' \times_{\mathbb{C}^\times} X$. The morphism $u : X_{\mathcal{Q}'} \to D$ has compact fibers, so the closure $\overline{j(C)}$ of the image of j is also a finite type curve over D. $\overline{j(C)}$ is not necessarily prestable. However, by using resolution of singularities, we may obtain a prestable curve Y'' (with a resolution map $r : Y'' \to \overline{j(C)}$); base change may also be required at this step. This gives us a sequence of maps (over D)

$$\begin{tikzcd}
Y'' \arrow[r, "r"] & \overline{j(C)} \arrow[r, hook, "j"] & X_{\mathcal{Q}'} \arrow[r, "pr"] & Y' \arrow[r, "c"] & Z \text{ ,}
\end{tikzcd}$$

where $pr : X_{\mathfrak{c}_0^*\mathcal{P}} \to \Sigma_0'$ is the bundle structure map. The composition $c_r = \mathfrak{c}_0 \circ pr \circ j \circ r : Y'' \to Z$ is necessarily a contraction map. We denote the pullback $c_r^*\mathcal{R}$ by $\mathcal{Q}''$. We denote the lifts of the marked points z_i by y_i''.

Pulling back $\mathcal{R}$ step by step from C to Y'', we get a sequence of bundles, the last of which is $c_r^*\mathcal{P}$, as in the diagram below.

$$\begin{tikzcd}
\mathcal{Q}'' \arrow[rr] \arrow[d] & & \mathcal{Q}' \times X \arrow[r, "pr_1"] \arrow[d] & \mathcal{Q}' \arrow[r] \arrow[d] & \mathcal{P} \arrow[d] \\
Y'' \arrow[r, "r"] & \overline{j(C)} \arrow[r, hook, "j"] & X_{\mathcal{Q}'} \arrow[r, "pr"] & Y' \arrow[r, "c"] & Z
\end{tikzcd}$$

We also get a section $s' : \mathcal{Q}'' \to X$ from the composition

$$\mathcal{Q}'' \to \mathcal{Q}' \times X \to X.$$

The collection $(Y'', y_i'', \mathcal{Q}'', s')$ is a map to $[X/\mathbb{C}^\times]$, but not necessarily a Gieseker map, as the curve may have unstable components carrying a trivial bundle and a trivial section. We obtain the desired extension by contracting these unstable components.

Separability: Now suppose that we are given two different pairs $(Y_1, y_{1,i}, \mathcal{Q}_1, s_1)$ and $(Y_2, y_{2,i}, \mathcal{Q}_2, s_2)$ which both extend the given family over D compatibly with the given Giesker map $(Z, z_i, \mathcal{R})$ to $[\mathrm{pt}/\mathbb{C}^\times]$. We may freely suppose that both extensions are defined over the same base extension.

Consider the fiber product $Y_1 \times_Z Y_2$. Our assumptions imply that $Y_1|_{D^\times} = Y_2|_{D^\times}$ and that the special fibers of Y_1 and Y_2 both contract onto the special fiber of Z. It follows that all the maps in the bottom diamond of the diagram below are contraction maps.

$$\begin{tikzcd}
 & \mathcal{Q} \arrow[dl, "f"'] \arrow[d] \arrow[dr, "f_2"] & \\
\mathcal{Q}_1 \arrow[d] \arrow[dr] & Y_1 \times_Z Y_2 \arrow[dl] \arrow[dr] & \mathcal{Q}_2 \arrow[d] \arrow[dl] \\
Y_1 \arrow[dr] & \mathcal{R} \arrow[d] & Y_2 \arrow[dl] \\
 & Z &
\end{tikzcd}$$

Moreover the two sections $\mathcal{Q} \to \mathcal{Q}_1 \to X$ and $\mathcal{Q} \to \mathcal{Q}_2 \to X$ agree on the open dense set $\mathcal{Q}|_{D^\times}$. X is separated, so the two sections agree. The Gieseker map obtained by contracting any unstable components in $Y_1 \times_Z Y_2$ is unique, so it follows that the two given families are isomorphic.

Deligne-Mumford: (Sketch of proof) Let C_v be a component of C. If C_v is contracted to a point by the section-forgetting morphism F_s, then $\mathcal{P}|_{C_v}$ is trivial, so $s|_{C_v}$ must be equivalent to a non-trivial map $C_v \to X$. We know from Gromov-Witten theory that such maps admit only finitely many automorphisms.

On the other hand, if C_v is stable, then the existence of a non-trivial section on C can only reduce the number of automorphisms.

□

Now consider the stack $\mathfrak{M}_{g,I,\nu_*\beta}([\mathrm{pt}/\mathbb{C}^\times])$ the stack of all degree $ft_*\beta$ maps from prestable curves to $[\mathrm{pt}/\mathbb{C}^\times]$. This stack is smooth. We obtain a morphism

$$\widetilde{F}_s : \widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times]) \to \mathfrak{M}_{g,I,ft_*\beta}([\mathrm{pt}/\mathbb{C}^\times])$$

by forgetting the section s, but not contracting any unstable components.

Theorem 6.4. *$L_{\widetilde{F_s}}$ admits a relative perfect obstruction theory.*

Recall from [BF97] that a *relative perfect obstruction theory* for $L_{\widetilde{F_s}}$ is pair (E, e) consisting of an element E of the derived category of $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$, and a homomorphism $e : E \to L_{\widetilde{F_s}}$ in the derived category, such that

1. $E = [E^{-1} \to E^0]$ is locally equivalent to a two-term complex of locally free sheaves.
2. $H^0(e)$ is an isomorphism.
3. $H^{-1}(e)$ is a surjection.

Proof of Theorem 6.4. Our proof is an almost word-for-word copy of the one given by Behrend & Fantechi in [Beh97] and [BF97].

Fix a curve C and a principal $\mathbb{C}^\times$-bundle $p : \mathcal{P} \to C$, and let Γ denote the space $\mathrm{Hom}_{\mathbb{C}^\times}(\mathcal{P}, X)$ of sections. Γ comes equipped with "universal" families, illustrated below.

$$\begin{tikzcd}
\mathcal{P} \times \Gamma \arrow[r, "s"] \arrow[d, "p \times \mathrm{id}_\Gamma"] & X \arrow[d, "\rho"] \\
C \times \Gamma \arrow[r, "\phi_s"] \arrow[d, "\pi"] & {[X/\mathbb{C}^\times]} \\
\Gamma &
\end{tikzcd}$$

It follows from the functorial properties of the cotangent complex that we have a morphism $\tilde{e} : s^*L_X \to p^*\pi^*L_\Gamma$. If we take $\mathbb{C}^\times$-invariants in the pushdown via p, we get

$$\tilde{e}' : (p_*s^*L_X)^{\mathbb{C}^\times} \to \pi^*L_\Gamma.$$

Tensoring with the dualizing complex of C, we obtain a morphism

$$\tilde{e}'' : \omega_C \otimes (p_*s^*L_X)^{\mathbb{C}^\times} \to \omega_C \otimes \pi^*L_\Gamma = \pi^!L_\Gamma.$$

Then, by adjunction, we have a morphism

$$\tilde{e}'' : R\pi_*(\omega_C \otimes (p_*s^*L_X)^{\mathbb{C}^\times}) \to L_\Gamma.$$

Finally, it follows from Verdier duality that

$$R\pi_*(\omega_C \otimes (p_*s^*L_X)^{\mathbb{C}^\times}) = R\pi_*(p_*s^*T_X)^{\mathbb{C}^\times},$$

and so we have a morphism

$$e : [R\pi_*(p_*s^*T_X)^{\mathbb{C}^\times}]^\vee \to L_\Gamma.$$

This morphism is a perfect obstruction theory for L_Γ; the proof is more or less the same as in [BF97]. Moreover, all of the objects here generalize well to the relative case, and therefore apply to the universal family. Thus, we have a perfect relative obstruction theory

$$e : E = [R\pi_*(p_*s^*T_X)^{\mathbb{C}^\times}]^\vee \to L_{\widetilde{F_s}},$$

where now π, p, and s refer to the universal families on the moduli stack. $\square$

Given this perfect obstruction theory, we can define the *virtual structure sheaf* $\mathcal{O}^{vir} = \mathcal{O}^{vir}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])}$. This is an element of the bounded derived category of coherent sheaves on $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$, which may be thought of as a family of virtual fundamental K-homology cycles on the fibers of F_s. It is defined using the virtual normal cone machinery developed by Behrend & Fantechi [BF97]. The definition we give here seems to have first appeared in print in [Lee04].

First, recall the *intrinsic normal cone* [BF97]. This is a cone stack $\mathfrak{I}_{\mathcal{X}}$ associated canonically to a Deligne-Mumford stack $\mathcal{X}$. It is defined locally on an étale open set $U \to \mathcal{X}$ by choosing an embedding $\iota : U \to W$ of U into a smooth scheme W and then setting $\mathfrak{I}_{\mathcal{X}}|_U = [C_{U/W}/\iota^* T_W]$, where $C_{U/W}$ denotes the normal cone of U in W and T_W is the tangent bundle of W. This construction is independent of the choice of embedding and glues nicely to give $\mathfrak{I}_{\mathcal{X}}$. Moreover, the construction works in the relative case, giving a normal cone $\mathfrak{I}_f = \mathfrak{I}_{\mathcal{X}/\mathcal{Y}}$ for any Deligne-Mumford morphism $f : \mathcal{X} \to \mathcal{Y}$ of $\mathcal{X}$ to a smooth, unobstructed, equidimensional $\mathcal{Y}$.

We will denote the relative intrinsic normal cone of $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$ relative to $\mathfrak{M}_{g,I,ft_*\beta}(\mathrm{pt}/\mathbb{C}^\times)$ by $\mathfrak{I}_{\widetilde{F}_s}$. The existence of a perfect relative obstruction theory for $\widetilde{F}_s$ implies [BF97] that there exists a closed embedding

$$i : \mathfrak{I}_{\widetilde{F_s}} \to [E^1/E^0]$$

where the two-term complex $E^\vee = [E^0 \to E^1]$ of vector bundles is the dual of the complex E, and $[E^1/E^0]$ the quotient stack of E^1 by the action of E^0.

Definition 6.5 ([Lee04])**.** The *virtual structure sheaf* $\mathcal{O}^{vir}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])}$ is the element of the bounded derived category of coherent sheaves $D(\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times]))$ defined by the derived tensor product

$$\mathcal{O}^{vir}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])} = \mathcal{O}_{\mathfrak{I}_{F_s}} \overset{L}{\underset{[E^1/E^0]}{\bigotimes}} \mathcal{O}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])}.$$

F_s is proper and Deligne-Mumford, so there exists a pushforward along it:

$$(F_s)_* : K^0(\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])) \to K^0(\widetilde{\mathcal{M}}_{g,I,ft_*\beta}(\mathrm{pt}/\mathbb{C}^\times)).$$

But F_s is obstructed, so this pushforward does not have good properties. We correct this by using the *virtual pushforward*, defined by

$$(F_s)^{vir}_![V] = (F_s)_*[V \overset{L}{\underset{\mathcal{O}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])}}{\bigotimes}} \mathcal{O}^{vir}_{\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])}].$$

Thus, we have established the existence of virtual pushforwards along F_s. The next step in defining Gromov-Witten invariants for $[X/\mathbb{C}^\times]$ is to introduce a notion of "admissible class" on $\widetilde{\mathcal{M}}_{g,I,\beta}([X/\mathbb{C}^\times])$, and show that the virtual pushforward of such a class is an admissible class on $\widetilde{\mathcal{M}}_{g,I}(\mathrm{pt}/\mathbb{C}^\times)$. We intend to address this question in a future paper.

References

[AB83] M. F. Atiyah and R. Bott, *The Yang-Mills equations over Riemann surfaces*, Philos. Trans. Roy. Soc. London Ser. A **308**(1505), 523–615 (1983).

[AGV08] D. Abramovich, T. Graber and A. Vistoli, *Gromov-Witten theory of Deligne-Mumford stacks*, Amer. J. Math. **130**(5), 1337–1398 (2008).

[Ale04] V. Alexeev, *Compactified Jacobians and Torelli map*, Publ. Res. Inst. Math. Sci. **40**(4), 1241–1265 (2004).

[AV02] D. Abramovich and A. Vistoli, *Compactifying the space of stable maps*, J. Amer. Math. Soc. **15**(1), 27–75 (electronic) (2002).

[Beh97] K. Behrend, *Gromov-Witten invariants in algebraic geometry*, Invent. Math. **127**(3), 601–617 (1997).

[BF97] K. Behrend and B. Fantechi, *The intrinsic normal cone*, Invent. Math. **128**(1), 45–88 (1997).

[BM96] K. Behrend and Y. Manin, *Stacks of stable maps and Gromov-Witten invariants*, Duke Math. J. **85**(1), 1–60 (1996).

[Cap94] L. Caporaso, *A compactification of the universal Picard variety over the moduli space of stable curves*, J. Amer. Math. Soc. **7**(3), 589–660 (1994).

[Cap08] L. Caporaso, *Néron models and compactified Picard schemes over the moduli stack of stable curves*, Amer. J. Math. **130**(1), 1–47 (2008).

[CG07] T. Coates and A. Givental, *Quantum Riemann-Roch, Lefschetz and Serre*, Ann. of Math. (2) **165**(1), 15–53 (2007).

[CGMiRS02] K. Cieliebak, A. R. Gaio, I. Mundet i Riera and D. A. Salamon, *The symplectic vortex equations and invariants of Hamiltonian group actions*, J. Symplectic Geom. **1**(3), 543–645 (2002).

[CGS00] K. Cieliebak, A. R. Gaio and D. A. Salamon, *J-holomorphic curves, moment maps, and invariants of Hamiltonian group actions*, Internat. Math. Res. Notices **16**, 831–882 (2000).

[CR02] W. Chen and Y. Ruan, Orbifold Gromov-Witten theory, in *Orbifolds in mathematics and physics (Madison, WI, 2001)*, volume 310 of *Contemp. Math.*, pages 25–85, Amer. Math. Soc., Providence, RI, 2002.

[FLN08] E. Frenkel, A. Losev and N. Nekrasov, *Instantons beyond topological theory II*, (2008), arXiv:hep-th/0803.3302.

[Gie84] D. Gieseker, *A degeneration of the moduli space of stable bundles*, J. Differential Geom. **19**(1), 173–206 (1984).

[GW08] E. Gonzalez and C. Woodward, *Area dependence in gauged Gromov-Witten theory*, (2008), arXiv:math/0811.3358.

[GW09] E. Gonzalez and C. Woodward, *Gauged Gromov-Witten theory for small spheres*, (2009), arXiv/0907.3869.

[Har77] R. Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.

[HM98] J. Harris and I. Morrison, *Moduli of curves*, volume 187 of *Graduate Texts in Mathematics*, Springer-Verlag, New York, 1998.

[Kle84] S. L. Kleiman, The structure of the compactified Jacobian: a review and announcement, in *Geometry seminars, 1982–1983 (Bologna, 1982/1983)*, pages 81–92, Univ. Stud. Bologna, Bologna, 1984.

[Lee04] Y.-P. Lee, *Quantum K-theory. I. Foundations*, Duke Math. J. **121**(3), 389–424 (2004).

[Mar81] M. Maruyama, *On boundedness of families of torsion free sheaves*, J. Math. Kyoto Univ. **21**(4), 673–701 (1981).

[MiR00] I. Mundet i Riera, *A Hitchin-Kobayashi correspondence for Kähler fibrations*, J. Reine Angew. Math. **528**, 41–80 (2000).

[MiR03] I. Mundet i Riera, *Hamiltonian Gromov-Witten invariants*, Topology **42**(3), 525–553 (2003).

[MiRT04] I. Mundet i Riera and G. Tian, *A compactification of the moduli space of twisted holomorphic maps*, (2004), arXiv: math/0404407.

[NS97] D. S. Nagaraj and C. S. Seshadri, *Degenerations of the moduli spaces of vector bundles on curves. I*, Proc. Indian Acad. Sci. Math. Sci. **107**(2), 101–137 (1997).

[NS99] D. S. Nagaraj and C. S. Seshadri, *Degenerations of the moduli spaces of vector bundles on curves. II. Generalized Gieseker moduli spaces*, Proc. Indian Acad. Sci. Math. Sci. **109**(2), 165–201 (1999).

[Ses82] C. S. Seshadri, *Fibrés vectoriels sur les courbes algébriques*, volume 96 of *Astérisque*, Société Mathématique de France, Paris, 1982, Notes written by J.-M. Drezet from a course at the École Normale Supérieure, June 1980.

[Sim94] C. T. Simpson, *Moduli of representations of the fundamental group of a smooth projective variety. I*, Inst. Hautes Études Sci. Publ. Math. **79**, 47–129 (1994).

[Tel00] C. Teleman, *The quantization conjecture revisited*, Ann. of Math. (2) **152**(1), 1–43 (2000).

[TW03] C. Teleman and C. T. Woodward, *The Index Formula on the Moduli of G-bundles*, (2003), arXiv:math.AG/0312154.

[Tel05] C. Teleman, *Loop Groups & G-Bundles on Curves*, (2005) Lecture at the Summer Institute in Algebraic Geometry, Seattle.

[Wit88] E. Witten, *Topological Sigma Models*, Commun. Math. Phys. **118**, 411 (1988).

Department of Mathematics, University of California, Berkeley, CA 94720, USA

E-mail address: `frenkel@math.berkeley.edu`
E-mail address: `teleman@math.berkeley.edu`
E-mail address: `ajt@math.sunysb.edu`